

Table 1 Suppl. Description of PCR primer pairs for amplifying. The primers designed by <sup>1</sup> - Guzmán and Alvarez (2012), <sup>2</sup> - Shariflou and Sharp (1999), and <sup>3</sup> - Nakamura *et al.* (2002).

Primer	Sequence (5' → 3')		
Wx1Fw <sup>1</sup>	TTGCTGCAGGTAGCCACACC		
Wx2Fw <sup>1</sup>	ATGGTCATCTCCCCGCGCTA		
Wx2Rv <sup>1</sup>	GTTGACGGCGAGGAACTTGT		
Wx3Rv <sup>1</sup>	TTCTCTTTCAGGGAGCGGC		
SUN1R <sup>2</sup>	ATAGGCACAACCCCTAAC		
AFC <sup>3</sup>	TCGTGTTTCGTCGGCGCCGAGATGG		
AR2 <sup>3</sup>	CCGCGCTTGTAGCAGTGGAAGTACC		
BDFL <sup>3</sup>	CTGGCCTGCTACCTCAAGAGCAACT		
BRD <sup>3</sup>	CTGACGTCCATGCCGTTGACGA		
Wx-3°2Fw	CACCGGGAAGAAGAAGTTTG		

Pair	PCR conditions		
	initial denaturation	35 cycles	final extension
Wx3°2Fw / SUN1R		45 s at 94 °C 30 s at 64 °C 1 min 40 s at 72 °C	
Wx2Fw / Wx2Rv		30 s at 94 °C 30 s at 66 °C 1 min 30 s at 72 °C	
Wx1Fw / Wx3Rv	3 min at 94 °C	50 s at 94 °C 50 s at 66 °C 3 min at 72 °C	5 min at 72 °C
AFC / AR2		40 s at 94 °C	
BDFL / BRD		30 s at 66 °C 1 min 10 s at 72 °C	

Table 2 Suppl. The origin and characteristics of *Wx* alleles evaluated.

Genes	Cultivar	NCBI accession	DNA size [bp]		Deduced protein		pI
			complete	coding	size [aa]	Mr [Da]	
<i>Wx-A1a</i>	Chinese Spring	AB019622	2781	1815	605	66.556	7.42
<i>Wx-A1a'</i>	Blanqueta	KJ185387	2781	1815	605	66.434	7.81
<i>Wx-A1h</i>	Astrodur	KJ185385	2780	1814	-	-	-
<i>Wx-B1a</i>	Chinese Spring	AB019623	2794	1818	606	66.345	7.39
<i>Wx-B1e</i>	Astrodur	KJ185386	2793	1818	606	66.313	7.39
<i>Wx-B1e'</i>	Lobeiro oscuro/ruivo	KJ185388	2804	1815	605	66.227	7.39
<i>Wx-B1f</i>	Mourisco	KJ185389	2793	1818	606	66.313	7.39




Fig. 1 Suppl. A schematic representation of the central region of *Wx-B1* gene among cvs. Turkey Red, Astrodur, Lobeiro, and Eureka. Alleles are not indicated in the case of partial sequences. The grey bars represent sequences; exons are indicated with “E” (the yellow arrows) and introns with “I” (the yellow lines) below the sequences. Nucleotide differences are highlighted in black and InDel events by the *hyphens*.




Fig. 2 Suppl. A schematic representation of the complete *Wx* gene and variability found among alleles *Wx-B1a* (CS, Chinese Spring), *Wx-B1e* (Ast, Astrodur), *Wx-B1e'* (Lob, Lobeiro) and *Wx-B1f* (Mou, Mourisco) analysed in this study. The grey bars represent sequences; exons are indicated with “E” (the yellow arrows) and introns with “I” (the yellow lines) below the sequences. Nucleotide differences are highlighted in black and InDel events by the *hyphens*.