

Table 1 Suppl. Sequences of oligonucleotide primers used in this study. The 5' end of the primers was extended with 8 bp to generate a *Sph*I digest site in the forward primers and a *Bgl*II digest site in the reverse primers (*in bold italics*). *GUS* - β -glucuronidase, TAIL - thermal asymmetric interlaced.

Primer	Sequence	Usage
SP1	AGAGCAACTTCAGCATCGCAGAGGAC	TAIL PCR
SP2	CCTTCGCTTAGCAAACGTGACCTGAC	
SP3	GTTCAACTCGTCCTCTCCCCATTCT	
pPlacSEP1.1FL	GGCATGCC ACTAATATAAGCACAGAA	<i>GUS</i> fusion vectors construction
pPlacSEP1.1D1	AGCATGCC GTTCATTTTGCTTCCTTTG	
pPlacSEP1.1D2	GGCATGC ATGGGTTTAGCAATGAGTGT	
pPlacSEP1.1D3	TGCATGC ACATCAACCATTTCCCCAT	
pPlacSEP1.1D4	GGCATGCGG ACCTGTGGTTGTGGGGCT	
pPlacSEP1.1D5	CGCATGC ACCAGGGCGTTAAATG	
pPlacSEP1VR	TC AGATCT ACTCGTCCTCTCCCCAT	
BarnaseVF	CC AGATCT CAGGTTATCAACACGTT	
BarstarVR	CGGTGACC ATATTGTTTCATCTCCCAT	
35SVF	CTCTAGAC GTAAAGGGATGACGCACAAT	<i>Barnase-mi35S-Barstar</i> construction
35SVR	TCACCAGTCCCCCGTGTCTCTCC	
1303R	CTGAATGCCACAGGCCGTCGAGTT	plasmid verification transgenic plants verification
pPlacSEP1VF	GGCAGTGACATTGGCTCCATCTAAG	
GUSVR	TGTAGAGCATTACGCTGCGATGGATT	

Table 2 Suppl. Putative *cis*-elements identified in the *Platanus acirifolia* *SEPALLATA1* promoter sequence.

Motif	Sequence	Distance from ATG	Function
CAAT-box	CCAAT/CAAT/CAATT	-561,-569,-585,-868,-988, -1,149,-1,210,1,220,-1,390, -1,424,-1,813,-1,924,-1,998, -2,035, -2,085	common <i>cis</i> -acting element in promoter and enhancer regions
TATA-box	TATAAT/TATA	-4,-7,-320,-457,-548,-558,-592, -874,-1,130,-1,144,-1,331, -1,921,-1,981,	core promoter element for transcription start
A-box	CCGTCC	-435	<i>cis</i> -acting regulatory element
ACE	GACACGTTTG	-378	light responsive element
Box I	TTTCAA	-251	
GT1-motif	GGTTAA	-738,-772	
I-box	AAGATAAGA/GATAAG G	-277, -480	
GAG-motif	GGAGATG	-652,-1,684	
G-Box	CACGTT	-380	
IBOXCORE	GATAA	-30,-279,-482,-1,535	conserved sequence upstream of light-regulated genes
ARE	TGGTTT	-356	<i>cis</i> -acting regulatory element essential for the anaerobic induction
CGTCA-motif	CGTCA	-665	MeJA-responsiveness element
TCA-element	GAGAAGGATA/GAAGA AGAAG	-31, -40, -55, -58	salicylic acid responsiveness element
GAREAT	TAACAAR	-26,-1,108,-1,613	gibberellin-responsive element
GARE-motif	AAACAGA	-450, -826	
ARFAT	TGTCTC	-781	auxin responsive element
MBS	TAACTG	-313,-512	MYB binding site involved in drought response
HSE	AAAAAATATC	-263	heat stress responsiveness element
TC-rich repeats	AATTTCTCA	-1,284	<i>cis</i> -acting element involved in defense and stress responsiveness
GATABOX	GATA	-31,-280,-483,-1,040, -1,049,-1,536	tissue specific expression element
CCGTCC-box	CCGTCC	-435	meristem specific activation element
CARGCW8GAT	CTAATATAAG	-546	<i>cis</i> -acting element in promoter of MADS-box genes
POLLEN1LELAT52	AGAAA	-96,-208,-275,-411,-525, -538,-616,-622,-762,-795, -816,-971,-1,381,-1,526, -1,899	motif required for pollen specific expression in tomato
GTGANTG10	GTGA	-116,-925,-1,572	
Skn-1_motif	GTCAT	-664, -1694, -1847	<i>cis</i> -acting regulatory element required for endosperm expression
PROLAMINBOXOSTGCAAAG		-420	
GLUB1			
DOFCOREZM	AAAG	-24,-94,-173,-207,-225,-283,-367,-420,-427,-524,-619, -625,-680,-765,-792,-798, -1,082,-1,297,-1,378,-1,384,-1,510,-1,539, -1,843,-2,113	core site required for binding of endosperm specific proteins
SEF1MOTIF	ATATTTAWW	-1,124	soybean embryo factor
ACGTCBOX	GACGTC	-1,989	seed development relate element
UP1ATMSD	CCCATT	-2,010	regulate gene expression during initiation of axillary bud outgrowth
XYLAT	ACAAAGAA	-1,382	xylem development regulatory elements
LIBOXATPDF1	TAAATGYA	-1,852	trichome specific regulatory elements
WBOXATNPR1	TTGAC	-341,-2,063	negative regulatory elements
WBOXNTERF3	TGACY	-923,-1,458,-2,062	
WRKY71OS	TGAC	-115,-341,-924,-1,036, -1,459,-1,796,-2,063	