

REVIEW

Photosynthesis and antioxidative defense mechanisms in deciphering drought stress tolerance of crop plants

D. EASWAR RAO and K.V. CHAITANYA*

Department of Biotechnology, GIT, GITAM University, Visakhapatnam-530045, India

Abstract

Crop plants are regularly exposed to an array of abiotic and biotic stresses, among them drought stress is a major environmental factor that shows adverse effects on plant growth and productivity. Because of this these factors are considered as hazardous for crop production. Drought stress elicits a plethora of responses in plants resulting in strict amendments in physiological, biochemical, and molecular processes. Photosynthesis is the most fundamental physiological process affected by drought due to a reduction in the CO₂ assimilation rate and disruption of primary photosynthetic reactions and pigments. Drought also expedites the generation of reactive oxygen species (ROS), triggering a cascade of antioxidative defense mechanisms, and affects many other metabolic processes as well as affecting gene expression. Details of the drought stress-induced changes, particularly in crop plants, are discussed in this review, with the major points: 1) leaf water potentials and water use efficiency in plants under drought stress; 2) increased production of ROS under drought leading to oxidative stress in plants and the role of ROS as signaling molecules; 3) molecular responses that lead to the enhanced expression of stress-inducible genes; 4) the decrease in photosynthesis leading to the decreased amount of assimilates, growth, and yield; 5) the antioxidant defense mechanisms comprising of enzymatic and non-enzymatic antioxidants and the other protective mechanisms; 6) progress made in identifying the drought stress tolerance mechanisms; 7) the production of transgenic crop plants with enhanced tolerance to drought stress.

Additional key words: abiotic stresses, antioxidants, chlorophyll, net photosynthetic rate, osmolytes, PEPC, ROS, RuBPC, water use efficiency.

Introduction

Plants grow well only under an optimum range of environmental factors including radiation, temperature, air humidity, and water supply. Even a slight variation of these factors influences growth and productivity. Changes in the photoperiod, the amount of radiation and its spectral composition, nutrient affluence or starvation, drought or flooding, high-speed wind, UV radiation, anaerobic conditions, extreme temperatures, and air and

soil pollution, all these abiotic factors can cause stress on the plants (Versulues *et al.* 2006). Drought stress is one of the most vital, multidimensional abiotic stress factors that adversely affect plant growth, metabolism, and yield (Osakabe *et al.* 2014) and is responsible for decreasing the crop yield by up to 50 % (Zlatev and Lidon 2012). Drought stress is outlined by changes in water relations, physiological processes, alterations in the cell membrane

Submitted 8 August 2015, *last revision* 29 October 2015, *accepted* 11 November 2015.

Abbreviations: AA - ascorbic acid; ABA - abscisic acid; APX - ascorbate peroxidase; C2H2 - zinc finger domain; DREB - dehydration-responsive element binding; DRO - deeper rooting; GB - glycine betaine; GR - glutathione reductase; GRAS - gibberellic-acid insensitive repressor; GSH - glutathione; GSSG - glutathione disulfide; LEA - late embryogenesis abundant; MDA - monodehydroascorbate; MDHAR - monodehydroascorbate reductase; MYB - myeloblastosis; NADP-ME - nicotinamide adenine dinucleotide phosphate malic enzyme; NAR - net assimilation rate; NCED - 9-*cis*-epoxycarotenoid dioxygenase; NPR - natriuretic peptide receptor; PCR - pentose carbon reduction; PEPC - phosphoenolpyruvate carboxylase; PPK - phosphopyruvate dikinase; PS - photosystem; ROS - reactive oxygen species; RWC - relative water content; SOD - superoxide dismutase; SPS - sucrose phosphate synthase; STZ - salt tolerant zinc finger protein; WRKY - transcription factor with conserved sequence; WUE - water use efficiency; YUC - yucasin; ZAT - zinc transporter; ZFP - zinc finger proteins.

Acknowledgements: The research laboratory of K.V. Chaitanya is funded by grants from the Department of Biotechnology, Govt. of India (No. BT/PR14467/AGR/02/742/2010). Easwar Rao acknowledges DBT for the research fellowship.

* Corresponding author; Fax: +91-891-2790399, e-mail: viswanatha.chaitanya@gmail.com

structure, and ultra-structure of cell organelles (Yordanov *et al.* 2003). Plants experience drought stress because of two major reasons: due to a deficit water supply to the roots and high transpiration rates. These conditions

mostly prevail under arid and semi-arid climates (Rahdari and Hoseini 2012). To overcome the low water availability, plants adapt using stress avoidance and stress tolerance mechanisms (Lawlor 2013).

Plant growth and water relations under drought stress

A deeper root system, increased root to leaf surface, synthesis of new root tips, increased root length, and increased density are various adaptations of plant's root systems to decreased soil water content also help to increase hydraulic conductivity of the plants (Peña-Valdivia *et al.* 2010). Transport of water in plant roots is achieved either through the apoplastic pathway or the symplastic pathway; the latter of which includes plasma membrane or tonoplast, in which the transport is mediated by the expression of aquaporins (Giuliani *et al.* 2005). The up-regulation of aquaporin genes like *PIP1* and *RWC-3* helps the maize crop to increase water uptake capacity (Giuliani *et al.* 2005). Adaptations of plants to drought can be best understood by studying the genes and transcription factors responsible for root patterning. The transcription factor dehydration-responsive element binding *DREB1A* driven by a stress responsive promoter *rd29* from *Arabidopsis thaliana* helps in the root development of groundnut during drought stress (Shridhar *et al.* 2012). It has been reported that the *Arabidopsis AVP1* gene increases root growth during drought stress (Park *et al.* 2005). The *Deeper Rooting 1 (DRO1)* gene is responsible for the root depth increase due to the increased gravitropic response in rice root tips. This gene when introduced into IR64 genotype of rice by backcrossing resulted in an increased drought tolerance without any apparent loss in yield (Uga *et al.* 2013).

Plants undergo various changes to avoid water stress induced damage. Some changes are Morphological like increased development of root hairs, deepening of roots, and rolling of leaves. There are also physiological changes like alterations in carbon partitioning, osmotic adjustment, reduced stomatal conductance, and increased water use efficiency (WUE, Taji *et al.* 2004). Water use efficiency is usually calculated as net photosynthetic rate to transpiration rate ratio or also as biomass production to transpiration rate ratio (WUE_m). In the second case, it is influenced by the relative growth rate, leaf area ratio (LAR), and net assimilation rate (Van den Boogaard *et al.* 1997). Increased WUE is caused by a less reduction in photosynthetic rate than in transpiration rate due to stomatal closure (Chaves *et al.* 2009).

Annual crop plants adapt to drought stress by decreasing their leaf size. This is preceded by a delay in the flowering process (Sujata *et al.* 2013). Drought stress suppresses plant growth and development in crop plants. Water deficiency leads to a reduction in the dry matter of wheat (Ahmad *et al.* 2007). Accumulation of fresh mass in cowpea and common bean is inhibited more than

accumulation of dry mass (Augé *et al.* 2001) and so the dry mass/fresh mass ratio in the common bean increases (Ramos *et al.* 1999). Relative growth rate of durum wheat and bean plants decrease during drought stress by 25 % due to a reduction in the net assimilation rate (NAR) and changes in the photosynthetic rate. This is due to a decreased LAR (Lutts *et al.* 2004, Berova and Zlatev 2002).

Drought stress is one of the critical constraints for the growth and production of crop plants. Improvement of drought stress tolerance is dependent on primary factors like osmotic adjustment (Praba *et al.* 2009). Osmotic adjustment is one of the important components of drought resistance in crop plants, this involves accumulation of solutes and thus a decreased osmotic potential as well as increased pressure potential (Martinez *et al.* 2007). Osmotic adjustment helps the plants in maintaining a high relative water content (RWC) at a low leaf water potential, thereby sustaining the growth (Farooq *et al.* 2009). Osmotic adjustment has been observed in the leaves of many crops such as wheat (Budak *et al.* 2013), maize (Pei *et al.* 2010), sorghum (Assefa *et al.* 2010), chickpea (Krouma 2010), field pea (Zlatev and Lidon 2012), sunflower (Rauf and Sadaqat 2008), common bean (Guler *et al.* 2012), rice (Bunnag and Pongthai 2013), barley (Witcombe 2008), and pearl millet (Boyer *et al.* 2008).

Drought stress is associated with changes in the content of phytohormones (Yang *et al.* 2002). Cytokinins and abscisic acid (ABA) are important hormones that play a major role in drought tolerance (Raghavendra *et al.* 2010). Elevated ABA content in response to drought stress leads to many physiological changes like stomatal closure to avoid water loss through transpiration. Abscisic acid sensing and signalling are thought to be mediated according to the current model established in *Arabidopsis thaliana*, which involves: pyrabactin resistance 1 (PYR)/PYR1-like (PYL)/regulatory components of the ABA receptor (RCAR), protein phosphatase 2C (PP2C), and sucrose non-fermenting-1 (SNF1)-related protein kinase 2 (SnRK2) (Yoshida *et al.* 2015). Abscisic acid plays a role in protecting plants during drought depending on the crop, drought stress profile, and season (Blum 2015). Drought stress results in leaf senescence which is related to the decline in cytokinin content in various plant species. The *ipt* gene of cytokinin biosynthesis is reported to delay leaf senescence and promote stress resistance in several plant species (Merewitz *et al.* 2011).

Oxidative stress in plants

Production of reactive oxygen species (ROS): One of the most striking and deleterious results of the abiotic stress in crop plants is the production of toxic ROS in

different cellular and sub-cellular compartments (Vaahtera and Brosché 2011). Some ROS are considered to be the most potent reactive ions known.

Table 1. Localization and functions of ROS in plants.

ROS	Localization	Function
Singlet oxygen ($^1\text{O}_2$)	chloroplasts	mediates retrograde signalling through activation of second messengers
Superoxide radical ($\text{O}_2^{\cdot-}$)	ETC, mitochondria, chloroplasts, peroxisomes, cytosol	dismutated to H_2O_2
Hydrogen peroxide (H_2O_2)	peroxisomes, mitochondria, chloroplasts	oxidizes proteins; reacts with $\text{O}_2^{\cdot-}$ in a Fe catalysed reaction to form $\text{OH}^{\cdot}$
Hydroxyl radical ($\text{OH}^{\cdot}$)	chloroplasts	originates by Haber-Weiss/Fenton reaction; reacts with all biomolecules

They are generated due to the decreased content of intracellular CO_2 , this results in the transfer of electrons from the electron transport chain to oxygen at photosystem I (Reddy *et al.* 2004). Synthesis and production of ROS, such as superoxide radical ($\text{O}_2^{\cdot-}$), hydroxyl radical ($\text{OH}^{\cdot}$), hydroperoxyl radical ($\text{HO}_2^{\cdot}$), hydrogen peroxide (H_2O_2), alkoxy radical ($\text{RO}^{\cdot}$), peroxy radical ($\text{ROO}^{\cdot}$), singlet oxygen ($^1\text{O}_2$), and excited carbonyl (RO^*) leads to the damage of lipids, nucleic acids, and sugars (Vellosillo *et al.* 2010). Reactive oxygen species can damage virtually all macromolecules. This results in a severe damage to cellular components causing DNA lesions. This causes mutations that can often lead to irreparable metabolic dysfunction and cell death (Karuppanapandian *et al.* 2011). The ROS also act as regulators of cell division and differentiation in growing tissues (Schippers *et al.* 2012). The ROS are the by-products of different metabolic pathways and they are produced not only in chloroplasts but also in mitochondria, cell wall, peroxisomes, endoplasmic reticulum, and plasma membrane. During abiotic stresses plants undergo several mechanisms to combat increased ROS production, and in some plants ROS are purposefully generated as signalling molecules to control programmed cell death, pathogen defence, or stomatal closure (Apel and Hirt 2004). The expression of different transcription factors is enhanced by ROS and includes members of the transcription: WRKY transcription factor (consisting of a 60 amino acid domain with conserved amino acid sequence WRKYGQK at its N-terminal end, zinc transporter (ZAT), related to ABI3/VP1 (RAV), GRAS (plant-specific gene family of putative transcription factors whose name derived from its first three identified members, gibberellic acid insensitive (GAI), repressor of GAI (RGA), and scarecrow (SCR) (Pang and Wang 2010).

Singlet oxygen ($^1\text{O}_2$) is produced by the input of energy to the relatively stable oxygen resulting in removal of spin restriction and rearrangement of electrons. Two forms of $^1\text{O}_2$ can be generated according

to the access of one electron into the free orbitals. In both forms of $^1\text{O}_2$, the spin restriction is eliminated resulting in the increase of oxidizing ability. The chlorophyll pigments associated with the electron transport system are the primary sources of $^1\text{O}_2$. Singlet oxygen may also arise as by-product of lipoxygenase activity (Armin *et al.* 2014). Hydroxyl radicals ($\text{OH}^{\cdot}$) and $^1\text{O}_2$ are the most reactive molecular species. The hydroxyl radicals are generated in Fenton reactions in the presence of a suitable chelating agent like iron under a specific temperature and neutral pH conditions. Hydroxyl radicals can even be produced from $\text{O}_2^{\cdot-}$ and H_2O_2 (Keles and Unyayar 2004). The ROS are also produced during normal cellular metabolisms, but during drought stress the balance between the ROS production and elimination gets disturbed. The enormous ROS that are produced in young leaf cells during drought stress are eliminated by complex non-enzymatic and enzymatic antioxidant systems.

Role of ROS as signalling molecules: Apart from being the damaging agents, ROS also act as signalling molecules in promoting the adaptation of plants to abiotic stress (Foyer and Noctor 2000, Jaspers and Kangasjärvi 2010). Reactive oxygen species signalling is highly integrated with several plant hormone responses such as ABA induced stomata closure; auxin induced root gravitropism; gibberellic acid induced seed germination and hypersensitive responses; jasmonic acid induced programmed cell death; and salicylic acid induced lignin biosynthesis (Sharma *et al.* 2014). However, to date there is no clear understanding about the mechanism of ROS signalling and the perceiving of ROS signals prior to drought stress adaptation remains elusive. Under biotic stress ROS along with salicylic acid and methyl salicylate; jasmonic acid and methyl jasmonate; azelaic acid; auxin; ethylene; ABA, *etc.*, mediate systemic signals in plants to achieve systemic acquired acclimation (Mittler and Blumwald 2015). Most of the ROS generated by NADPH oxidases act as signaling molecules (Jaspers and Kangasjärvi 2010). Perceiving of

extracellular ROS might be done in multiple mechanisms with the possibilities of a protein redox regulation leading to ROS sensing or the oxidation of cell wall components or membranes that in turn is sensed by receptors to transduce the signal into the cytosol (Hancock *et al.* 2006). The intracellular redox status is sensed also by transcription factors such as NPR1, whose monomerization permits the translocation of protein into the nucleus, thereby enhancing the defense response (Tada *et al.* 2008). Candidates for the recognition of ROS in the apoplast include cysteine-rich protein receptor-like kinases (CRKs), small apoplastic proteins, and peptides

(Reddie and Carroll 2008). Cysteine-rich protein receptor-like kinases have been found to be transcriptionally regulated by oxidative stress, pathogen attack, and plant hormones. The small extracellular GRIM REAPER (GRI) protein is another candidate for ROS perception. A peptide derived from GRI can induce cell death in plants in the presence of a superoxide radical. However, existing evidence could support the probable role for CRKs and GRI in ROS perception; this hypothesis is yet to be experimentally tested (Wrzaczek *et al.* 2011).

Table 2. Expression of transcription factors in different plants and their functions in stress response.

Gene	Plant	Function
<i>DREB1C</i>	<i>Arabidopsis</i>	enhances desiccation tolerance
<i>ABF3</i>	rice	inhibits wilting and rolling
<i>DREB1A</i>	wheat, rice	delays wilting, improves drought tolerance
<i>AREB1</i>	<i>Arabidopsis</i>	enhances dehydration tolerance
<i>HARDY</i>	rice	induces high WUE and photosynthesis
<i>ABP9</i>	maize	increases photosynthesis under drought
<i>ABF1, ABF2</i>	rice	improves drought tolerance
<i>bZIP 23</i>	rice	improves drought tolerance
<i>NAC2</i>	rice	induces stomatal closure
<i>NAC6, NAC10</i>	rice	protects against oxidative stress
<i>SIAREB</i>	tomato	improves drought tolerance

Molecular responses to drought stress

The physiological and biochemical responses that are induced during drought are the result of an assortment of genes with diverse functions (Yamaguchi-Shinozaki and Shinozaki 2005). These gene products function in response to drought stress resulting in tolerance at the cellular level. Drought stress triggers certain signals resulting in the activation of transcriptional factors (TFs) involved in drought tolerance. These transcriptional factors include dehydration responsive transcription factors (DREBs) and C-repeat binding factors (CBFs) that bind to the dehydration response element (DRE), C-repeat terminal (CRT) *cis*-acting elements, ethylene responsive element binding factor (ERF), zinc-finger family, WRKY family, basic helix-loop-helix (bHLH) family, basic-domain leucine zipper (bZIP) family, NAC family, and homeodomain transcription family. Drought stress-induced gene expression was seen to be regulated by TFs belonging to bZIP, AP2/ERF, HD-ZIP, MYB, bHLH, NAC, NF-Y, EAR, and ZPT2 families (Rao *et al.* 2015). The consequences of drought stress include variation in the expression of late embryogenesis abundant (*LEA*) genes, leading after this to the synthesis of molecular chaperones that assist to conserve other proteins from degradation (Xoconostle-Cazares *et al.* 2011). The transfer of stress-inducible genes like the 9-*cis*-epoxycarotenoid dioxygenase (*NCED*) gene *via* gene transfer produces stress tolerant plants (Zhang *et al.*

2008b). Microarray analysis of stress-inducible genes from different plant species assist in investigating the functions of these genes to have knowledge of molecular mechanisms governing the drought stress tolerance in plants through genetic manipulation (Table 2).

Genes like *DREB2A*, *RD29B*, and *LEAs* are induced in mature tissues of plants subjected to mild osmotic stress (Skirycz *et al.* 2011). Recent studies reported that *Arabidopsis* plants overexpressing *yucasin 6* (*YUC6*), belonging to the flavin monooxygenases (FMOs) family, exhibits improved drought stress tolerance due to a low rate of water loss and controlled ROS accumulation (Cha *et al.* 2015). During drought, ABA is produced, which not only causes stomatal closure but also activates stress-related genes encoding protein kinases or phosphatases. These genes contain a conserved ABA-responsive element (ABRE)/CRT (C-repeat) that is characterized with the promoter region of *RD29A*. With the advent of environmental stress in plants, the *cis*-element *trans*-factors CBF/DREB1 and DREB2 are expressed and mediate the up-regulation of genes involved in drought tolerance (Uno *et al.* 2000). Several *cis*-acting basic leucine zipper (bZIP) transcription factors AREB/ABF can bind to ABRE, thus activating ABA-dependent gene expression. The reduced activity of ABA in ABA-deficient *aba2*, ABA-insensitive *abil*, as well as enhanced activity in the ABA hypersensitive *eral*

Arabidopsis mutants resulted in the different activation of these transcription factors. The ABA-dependent gene expressions depend upon a drought-inducible *RD22* gene. The *RD22* gene is activated by the binding of the MYC transcription factor AtMYC2 (RD22BP1) and MYB transcription factor AtMYB2 with the *cis*-elements of *RD22* promoter region, resulting in improved drought tolerance in transgenic plants (Abe *et al.* 2003). The drought inducible gene with NAC transcription factor *RD26* is up-regulated with ABA in transgenics and repressed in the *RD26* repressor lines. The transcription factors NAC016, NAP, and AREB1 function in the drought stress response in addition to affecting leaf senescence in *Arabidopsis* (Sakuraba *et al.* 2015).

The products of drought-inducible genes identified through microarray analysis in *Arabidopsis* are classified into two groups. The first group includes effector proteins: LEA, osmotin, chaperones, antifreeze proteins, mRNA binding proteins, water channel proteins, sugar and proline transporters, detoxification enzymes, and proteases. The second group includes regulatory proteins: proteins that regulate signal transduction, stress-responsive gene expression, transcription factors, protein kinases, protein phosphatases, enzymes of phospholipid metabolism, and signalling molecules. Ethylene response factors 5 and 6 activate a plethora of osmotic stress tolerance genes such as WRKY33, MYB51, and salt tolerance zinc finger protein (STZ) (Dubois *et al.* 2013) and they were also shown to provide resistance to oxidative stress (Wang *et al.* 2013) and biotic stress (Sun *et al.* 2010).

LEA proteins are extremely hydrophilic proteins, which were first described in cotton as accumulating in late phases of seed development, whose function correlates with the membrane stability; it also possesses resistance to desiccation, thus acting as important source of cellular dehydration tolerance in plants (Sivamani *et al.* 2000). The desiccation sensitive *Lindernia subracemosa* plants under drought stress up-regulate the synthesis of LEA proteins similar to those present in the resurrection plants (Bartels *et al.* 2007). The LEA proteins are encoded by *RD* (responsive to dehydration), *ERD* (early responsive to dehydration), *COR* (cold regulated), *KIN* (cold inducible), and *RAB* (responsive to

abscisic acid) genes that are induced during drought stress in different plant species (Shinozaki and Yamaguchi-Shinozaki 2000, Zhu 2002). The LEA proteins are found in a wide range of crop plants like *Gossypium hirsutum* (Galau *et al.* 1993), *Glycine max* (Shih *et al.* 2004), *Oryza sativa* (Moons *et al.* 1997), *Pisum sativum* (Grelet *et al.* 2005), and *Raphanus sativus* (Raynal *et al.* 1990) in response to drought stress (Table 3).

Osmotin is a 24-kDa protein and belongs to dehydrins (LEA2 proteins) that play an important role in stress tolerance mechanisms in plants. Transgenic plants with the over-expressing *OSMOTIN* gene accumulate a significantly more protein under drought stress, determining the role of osmotin in imparting tolerance to drought stress (Barthakur *et al.* 2001). Induction of osmotin transcription is induced by ABA at low leaf water potentials and the amount of osmotin transcript increases with increasing endogenous ABA content (Singh *et al.* 1989). Transcription of osmotin is inhibited by a plethora of tissue specific and hormonal signals that show similarity with ABRE responsive elements; this is associated with osmotic stress tolerance in plants (Raghothama *et al.* 1993). In tomato, water stress causes a substantial rise in the content of osmotin mRNA and protein (Grillo *et al.* 1995). Besides osmotic adjustments, osmotin also safeguards the native structure of proteins during drought stress and is also involved in repairing denatured proteins (Goel *et al.* 2010). Transgenic cotton plants that have been transformed with the tobacco *OSMOTIN* gene have an increased tolerance to osmotic stress (Parkhi *et al.* 2009).

Aquaporins are channel proteins localized in the plasma membrane and intracellular membranes and they facilitate the transport of water, ions, and gases (Maurel *et al.* 2008). The major intrinsic proteins (MIP) family includes aquaporins that control water movement through cells and maintain plant water relations especially during drought in which the water supply is decreased (Šurbanovski *et al.* 2013). Aquaporins regulate water conductance at key “gatekeeper” cell layers; this causes a change in the whole plant water flow and plant water potential (Chaumont *et al.* 2014). Functions of aquaporins are reported to be involved in root hydraulics

Table 3. Expression of genes that encodes polypeptides related to LEA proteins in different plants during drought stress.

Gene	Plant	Related LEA proteins
<i>Em</i>	wheat	D19-LEA-protein (group 1)
<i>pRAB16A</i>	rice	D11-LEA-protein (group 2)
<i>DREB1A</i>	maize	D11-LEA-protein (group 2)
<i>ERD10;ERD14</i>	<i>Arabidopsis</i>	D11-LEA-protein (group 2)
<i>Ha ds10</i>	sunflower	D19-LEA-protein (group 1)
<i>HVA1</i>	barley	D19-LEA-protein (group 1)
<i>HVA1</i>	wheat	D7-LEA-protein (group 3)
<i>HVA1</i>	barley	D11-LEA-protein (group 2)
<i>pLE4</i>	tomato	D11-LEA-protein (group 2)
<i>pcECP40</i>	carrot	D11-LEA-protein (group 2)

and in water transport from xylem to stomata, there by facilitating the plant adaptation to short term changes in soil moisture (Gomes *et al.* 2009). The plasmalemma and tonoplast are rich in subfamilies of aquaporins called plasma membrane intrinsic proteins (PIP) and tonoplast intrinsic proteins (TIP), respectively. The PIPs and TIPs mediate the central pathways for transport of water in the cells and maintains intracellular and transcellular water relations during stress. The PIPs are involved in numerous processes in response to drought stress. Drought stress in grape vine causes an increased expression of five *PIP* genes (Galmes *et al.* 2007). However, transcription of the *Arabidopsis AtPIP* aquaporin gene in roots undergo down-regulation under drought stress preventing water back flow from the root to drying soil (Alexandersson *et al.* 2005). But it remains elusive whether the decreased expression of aquaporins

during drought is a deliberate or spontaneous phenomenon.

Zinc finger proteins (ZFP) play a crucial role in stress adaptations through their DNA binding or RNA binding interactions. Zinc finger domain (C2H2) families of ZFPs are strongly induced in plants under osmotic stress (Sakamoto *et al.* 2004). About 134 C2H2 ZFPs are reported in the *Arabidopsis* genome. Over-expression of rice *OsZFP252* augments the tolerance of rice seedlings to drought, which is shown by a remarkable survival rate of 79 % (Xu *et al.* 2008). The WRKY transcription factors are involved in the regulation of plant growth and development, and confer tolerance to biotic stresses (Ulker and Somssich 2004). Expression of *OsWRKY11* in rice under the *HSP 101* promoter provides tolerance to drought (Wu *et al.* 2009).

Photosynthesis during drought

Photosynthesis is an important physiological process occurring in all green plants (Ashraf and Harris 2013). The rate of photosynthesis is adversely affected by drought stress due to stomatal and non-stomatal limitations (Kawamitsu *et al.* 2000, Yordanov *et al.* 2003, Samarah *et al.* 2009). A decrease in stomatal conductance with a decrease in leaf water potential results in a reduced net photosynthetic rate and so a decreased assimilate production resulting in a reduced growth and yield (Nikinmaa *et al.* 2013). Stomatal closure is supposed to be one of the main reasons for the decreased net photosynthesis rate (Boyer 1970, Flexas *et al.* 2004). Stomatal closure leads to a decrease in intracellular CO₂ concentrations, which in turn affects Rubisco as well as other enzymes such as sucrose phosphate synthase (SPS) and nitrate reductase. Specific changes in membrane potential and cytosolic pH play an important role in determining the direction and capacity of ion transport in guard and subsidiary cells (Mumm *et al.* 2011). During drought, the activities of different C₄ photosynthetic enzymes like NADP malic enzyme (NADP-ME), phosphopyruvate dikinase (PPDK), and fructose-1,6-bisphosphatase decrease significantly, suggesting the relevance of these enzymes for drought responses (Du *et al.* 1998).

The ability of plants to acclimate to different abiotic stresses varies from species to species depending on the carbon fixation pathways (C₃, C₄, and CAM). About 85 % of higher plants represent C₃, 5 % are C₄, and the rest 10 % belongs to CAM. Drought stress represses cell growth and photosynthesis, and also affects the whole metabolism by adversely affecting the activities of different photosynthetic enzymes. The structural and biochemical impairment of light-dependent reactions as well as carboxylation processes of photosynthesis are the consequences of severe drought stress (Ghotbi-Ravandi *et al.* 2014).

The decrease in the activity of Calvin cycle enzymes

has also been observed in drought stressed plants (Maroco *et al.* 2002). Prolonged drought stress decreases the RuBP content due to the inadequate supply of ATP or NADPH to pentose carbon reduction (PCR) cycle. Water deficit affects Rubisco activase which regulates the conformational structure of the Rubisco active site by releasing tight binding inhibitors at the active site utilising ATP (Athar *et al.* 2005). In C₄ plants, fixation of CO₂ in the cytosol of mesophyll cells is mediated by phosphoenolpyruvate carboxylase (PEPC) to form oxaloacetate. Drought induces the accumulation of PEPC transcripts and PEPC proteins as well as an increased activity of PEPC. NADP-malic enzyme (NADP-ME) catalyses the oxidative decarboxylation of malate in the presence of NADP⁺ to produce pyruvate, CO₂, and NADP(H) (Doubnerová *et al.* 2011). The membrane damage caused during severe drought requires membrane lipid regeneration by synthesising fatty acids. The NADP-ME helps in providing the necessary NADH and pyruvate for the synthesis of fatty acids. The NADPH required for the antioxidative defense systems, like the ascorbate-glutathione cycle involved in plant protection against drought stress, is also produced by NADP-ME (Hernández *et al.* 2012). The up-regulation of NADP-ME during drought stress also helps in modulating organic anionic metabolism in guard cells and in control of stomatal closure by malate degradation during day (Guo *et al.* 2009). The PPDK is found abundantly in C₄ plants and in low quantities in C₃ plants (seeds and mid veins); it helps in catalysing the reversible reaction converting ATP, Pi, phosphoenolpyruvate (PEP), AMP, and PPi. The cytosolic PPDK plays an important role in transport of amino acids during natural leaf senescence and in accelerating the nitrogen metabolism from leaves, thereby increasing the nitrogen content of seeds (Chastain *et al.* 2006, Taylor *et al.* 2010).

Photosystem (PS) II plays a key role in photosynthetic response to drought stress (Misra *et al.* 2012). Drought

stress causes a considerable damage to photosynthetic pigments, which leads to degradation of thylakoid membranes (Kannan *et al.* 2011). The effect of drought on photosynthesis is due to the change in chlorophyll content, by affecting chlorophyll components, and by

damaging the photosynthetic apparatus. Drought stress declines the content of chlorophylls in sunflower and the reduction of chlorophyll *b* is greater than of chlorophyll *a* (Jaleel *et al.* 2009).

Antioxidants

The production of ROS is the outcome of a plant metabolism that needs to be controlled to safeguard its cellular components (Saeidnejad *et al.* 2015). The natural antioxidants maintain the cellular redox balance by binding and inactivating the free radicals, which is crucial for the survival of plants during environmental stresses (Most and Jutta 2015). The antioxidant defense system comprising of the co-ordinated action of enzymatic and non-enzymatic antioxidants provide an efficient mechanism of controlling the toxicity rendered by ROS.

Non-enzymatic antioxidants include ascorbic acid (AA), glutathione (GSH), α -tocopherol, and carotenoids among which AA is one of the most important antioxidants synthesised in the majority of plant cell types, organelles, and apoplast (Smirnoff and Wheeler 2000, Giovannoni 2007). The non-enzymatic antioxidants play a vital role in sustaining photosynthetic efficiency and yield stability under severe stress conditions (Gupta and Thind 2015). Except in dry seeds, AA occurs in almost all tissues of plants. Ascorbate can readily donate an electron forming a monodehydroascorbate (MDHA) radical which on further oxidation yields an uncharged radical dehydroascorbate (DHA). The electron donating capability and formation of less reactive MDHA forms the basis for its antioxidant and scavenging activity (Venkataraman *et al.* 2004). Ascorbate reacts with a hydroxyl radical, superoxide, and singlet oxygen. Ascorbic acid provides membrane protection by directly scavenging an $\text{OH}^\cdot$ ion and regenerating tocopherol from the tocopheroxy radical. Ascorbate is also an important co-substrate of key antioxidant enzymes such as ascorbate peroxidase (APX) that detoxifies H_2O_2 , and 2-oxoacid dependent dioxygenases involved in the biosynthesis of ABA and gibberellic acid (Brossa *et al.* 2011). The role of ethylene and gibberellins in growth regulation has been studied in the leaf tissues of *Brachypodium distachyon* and maize subjected to a mild drought at different developmental stages (Verelst *et al.* 2013). Ascorbic acid is involved in plant cell growth and division, then it is also a co-factor for many enzymes, *e.g.*, AA acts as co-factor in the reaction where APX catalyses the reduction of H_2O_2 to water with concomitant production of MDHA that further yields DHA and AA under catalysis of monodehydroascorbate reductase (MDHAR, Asada 1999).

Glutathione (γ -glutamylcysteinglycine) is another major low molecular mass antioxidant present in plants and occurring in virtually all cellular components, such as chloroplasts, mitochondria, endoplasmic reticulum,

vacuoles, and cytosol, performing multiple functions. Glutathione is associated with the storage and transport of reduced sulphur (Kopriva *et al.* 2004), detoxification of xenobiotics *via* glutathione-S-transferase (Dixon *et al.* 2002), synthesis of phytochelatins, and in the regulation of redox homeostasis of the cell together with its oxidised form (GSSG) (Cobbett 2000). The central nucleophilic cysteine residue of GSH helps in scavenging cytotoxic H_2O_2 and reacts enzymatically with various other ROS. Glutathione plays a vital role in antioxidative defense by regenerating AA *via* the AA-GSH cycle (Halliwell 2006). Glutathione is precursor for synthesising phytochelatins; this is crucial for controlling cellular heavy metal content (Koffler *et al.* 2014).

Enzymatic antioxidants: Superoxide dismutases (SODs) belong to the family of metallo-enzymes that catalyse the disproportion of superoxide ($\text{O}_2^\cdot$) to O_2 and H_2O_2 . All aerobic organisms that are prone to oxidative stress require SODs to dismutate $\text{O}_2^\cdot$ yielding H_2O_2 . Based on the metal co-factor, SODs are classified into three types: Fe-SOD (localized in chloroplasts), Mn-SOD (localized in mitochondria), and Cu/Zn SOD (localized in chloroplasts, peroxisomes, and cytosol). These three isoenzymes show different sensitivity to H_2O_2 . The first plant SOD gene was cloned from maize (Cannon *et al.* 1987). Numerous SOD isoenzymes have been reported from plants like nine in maize (Baum *et al.* 1979), seven in *Arabidopsis* (Kliebenstein *et al.* 1999), and two Cu/Zn SODs in *Nelumbo nucifera* (Dong *et al.* 2011). Several studies indicated that an increased expression of the *MnSOD* gene enhances tolerance to many environmental stresses. The over-expression of *Cu/ZnSOD* enhances the tolerance of potato to drought stress (Kumar *et al.* 2013). The Fe-SOD has been detected only in chloroplasts of limited species and has an inconsistent distribution, but genome analysis in higher plants indicates the wide distribution of chloroplastic *Fe-SOD* genes (Kanematsu and Asada 1994). The role of SOD activity varies with different abiotic stress conditions depending on the distribution as well as the mechanism involved in overcoming stress (Karuppanapandian *et al.* 2011).

Catalase is a tetrameric heme containing enzyme that occurs in all aerobic eukaryotes; it catalyses the dismutation of H_2O_2 generated in peroxisomes into H_2O and O_2 (Moloudi *et al.* 2013). Catalase is a critical enzyme for maintaining the redox balance during oxidative stress. Three isoforms of the catalase enzyme are encoded by *Arabidopsis* genes *CAT-1*, *CAT-2*, and *CAT-3* which are distributed on separate chromosomes

and differentially expressed as well as independently regulated. The *CAT* genes are sensitive to numerous abiotic stress signals that regulate their expression (Dat *et al.* 2000). The damage of tissue metabolism in response to stress is overcome by dismutating toxic H_2O_2 due to increased catalase activity.

Ascorbate peroxidase is localised in chloroplasts, cytosol, vacuoles, and extracellular components; it detoxifies H_2O_2 to form H_2O and monodehydroascorbate, using ascorbate as electron donor (Adriano *et al.* 2015). The APX is a key enzyme in the ascorbate-glutathione cycle also known as Halliwell-Asada pathway. It was reported that drought stress up-regulates cytosolic APX in alfalfa nodules (Naya *et al.* 2007). Thylakoid APX (tAPX) and stroma APX (sAPX) are the two isoforms of chloroplastic APX. Thylakoid APX binds to the vicinity of PS I, so that it detoxifies even small quantities of H_2O_2 in its production site (Miyake *et al.* 1993). The expressions of rice thylakoid *OsAPX8* and *OsAPX4* are significantly down-regulated during drought stress, which is due to the distinct responses of *APX* genes in different species and to different magnitudes of stress (Rosa *et al.* 2010). Transgenic plants over-expressing the *SbpAPX* gene show an enhanced drought stress tolerance (Singh *et al.* 2014). The SOD and APX, which are present in chloroplasts and exist in soluble as well as thylakoid-bound forms, expedite rapid scavenging $O_2^{\cdot-}$ generated at the membrane surface and immediately convert it to H_2O_2 (Flexas *et al.* 2004). The over-expression of a key enzyme of ABA biosynthesis 9-cis-epoxy-carotenoid dioxygenase gene (*SgNCED1*) in transgenic tobacco plants results in a higher accumulation of ABA and regulation of APX activity under drought stress (Zhang *et al.* 2008b). The over-expression of the *Solanum lycopersicum* thylakoid-bound APX gene (*StAPX*) in tobacco plants shows a higher APX activity under salinity and drought stress (Sun *et al.* 2010).

Glutathione reductase (GR) is an NAD(P)H-dependent enzyme that catalyses the reduction of GSSG to GSH and thus maintains the reduced pool of GSH. Glutathione reductase plays a vital role in drought tolerance in higher plants by scavenging radicals and safeguarding the -SH group via the GSH-disulphide exchange reaction. The rate limiting step of the Halliwell-Asada pathway is catalysed by GR. The maximal activity of GR is observed in photosynthetic tissues that contain chloroplastic isoforms that differ from other GR isoforms from cytosol, mitochondria, and peroxisomes. The GSH and GR are involved in the detoxification of H_2O_2 generated by the Mehler reaction. The onslaught of drought stress enhances GR activity resulting in the accumulation of GSH that acts as a potential antioxidant and radical scavenger. One cytosolic isoform (OsGR2)

and two chloroplastic isoforms (OsGR1 and OsGR3) of GR have been reported in rice (Rouhier *et al.* 2006, Monodehydroascorbate reductase (MDHAR) together with dehydroascorbate reductase (DHAR) take part in ascorbate glutathione cycle Over-expression of DHAR and MDHAR in transgenic tobacco and *Arabidopsis* induce an enhanced tolerance against osmotic stresses (Eltayeb *et al.* 2007).

Secondary metabolites that function as antioxidants during drought stress: Tocopherols and tocotrienols, in particular α -tocopherol, are synthesised by all plants and form essential components of biological membranes; they prevent auto-oxidation of lipids thus making it an effective free radical trap. Tocopherols are mainly located in chloroplast membranes and protect lipids and other membrane components of PS II (Igamberdiev *et al.* 2004). During drought, tocopherol and AA content significantly increase in rice (Ahmad *et al.* 2008). In higher plants, oxidative stress triggers the expression of genes that synthesise tocopherols (Wu *et al.* 2007). Tocopherols function in protection against photo-oxidative damage, repairment of oxidizing radicals, and stabilization of membrane structure by changing its fluidity.

Polyphenols or phenolic acids serve as powerful antioxidants because of their ability to eliminate radical ions and function as metal chelators. Phenolics, including flavonoids, tannins, hydroxycinnamate esters, and lignin are distinct secondary metabolites that are abundant in plant tissues (Grace and Logan 2000). Polyphenolic compounds and flavonoids have the capacity to amend peroxidation kinetics by customising lipid packaging in order to decrease the fluidity of membranes thus restraining peroxidation reactions (Arora *et al.* 2000). Flavonoids have been studied extensively for their antioxidant activities (Lambert and Elias 2010). They have been reported to interfere with the activities of enzymes involved in ROS generation, quenching free radicals, chelating transition metals, and rendering them redox inactive in the Fenton reaction (Nakabayashi *et al.* 2014). The conversion of polyunsaturated fatty acids to oxygen containing derivatives is catalysed by lipoxygenase and prostaglandin synthetase. Flavonoids act as potent inhibitors for these enzymes and inhibit the reaction.

Carotenoids are lipophilic organic compounds found in plastids where they play a role in the plant metabolism against oxidative stress in addition to their major role as photosynthetic pigments. Carotenoids react with lipid peroxidation products to terminate chain reactions preventing the formation of 1O_2 by depleting excess excitation energy through the xanthophyll cycle (Collins 2001).

Osmolytes

Plants accumulate organic solutes in cytoplasm, such as proline, glycine-betaine, glucose, fructose, mannitol,

inositol, valine, isoleucine, *etc.*, in response to drought stress. These compatible solutes do not inhibit normal

metabolic reactions and act as osmoprotectants under drought, a condition that mediates a change in external osmolarity and maintains osmotic balance and continuous water influx (Tarczynski *et al.* 1993). Osmolytes also increase the stability of macromolecules without interfering with their functional activities (Yancey *et al.* 1982). They also scavenge free radicals under severe drought stress (Padmavathi and Rao 2013).

Proline plays an important role in safeguarding cells from damage caused during drought stress. It scavenges ROS and acts as molecular chaperone stabilizing the structure of proteins (Szabados *et al.* 2010). The activities of Δ^1 -pyrroline-5-carboxylate (P5C) synthetase and P5C reductase help in increasing proline biosynthesis, whereby plants combat abiotic stresses (Hare and Cress 1997).

Glycine betaine (GB) is an amphoteric quaternary amine that acts as compatible solute in most plants during drought stress. It performs an important function in plants when subjected to abiotic stresses by mediating osmotic adjustments and protecting cells from damage (Chen *et al.* 2008). Plants usually contain a low amount of GB, whereas *Arabidopsis* and few crop plants like rice and tomato do not naturally accumulate GB (Fariduddin *et al.* 2013). Abiotic stresses enhance the accumulation of GB in sugar beet, barley, spinach, sorghum, and maize, which stabilizes the structures and activities of enzymes, thereby maintaining the integrity of membranes (Rhodes *et al.* 1993, Yang *et al.* 2003).

Fructans are polymers of branched fructose molecules that are synthesised in vacuoles and serve as reserve

saccharides. Fructans are highly water soluble and are synthesised at low temperatures (Livingston *et al.* 2009). During drought stress, fructans release hexoses thereby stabilise membranes and maintain osmotic adjustment in plants like barley and rye (Spollen and Nelson 1994, Olien and Clark 1995, Valluru and Van den Ende 2008). Fructosyltransferases help in the synthesis of fructans in plants; its introduction to fructan non-accumulating plants like tobacco and rice stimulates fructan production and enhances drought tolerance (Livingston *et al.* 2009).

Another important compatible osmolyte is trehalose, a non-reducing disaccharide that is present in low amounts in many plants but vary during environmental stresses. It helps in maintaining membrane stability and osmotic balance. Trehalose is synthesised in plants through the production of the intermediate trehalose-6-phosphate. Transgenic expressions of trehalose biosynthetic enzymes in various crop plants like rice (Jang *et al.* 2003), tomato (Cortina *et al.* 2005), and potato (Goddijn *et al.* 1997) result in trehalose accumulation leading to stress tolerance.

Mannitol is synthesised naturally in various plant species except a few like wheat. In celery, almost a half of fixed CO₂ is converted to mannitol, equal in proportion to that of sucrose. The low water potential of plants due to drought stress results in an increased accumulation of mannitol that is crucial for maintaining membrane stability. The over-expression of gene encoding mannitol-1-phosphate dehydrogenase results in the enhanced tolerance of *Arabidopsis* (Thomas *et al.* 1995) and tobacco (Tarczynski *et al.* 1992, 1993).

Transgenic plants with enhancing drought stress tolerance

Marked decreases in crop production due to abiotic stresses lead to a demand for the development of drought stress resistant crop plants obtained through transgenic approach (Vinocur *et al.* 2005). The induction of drought stress in plants triggers a plethora of genes which play a crucial role in the defence pathways, these were categorised into three groups: 1) those involved in the protection of proteins and membranes against damage incurred by ROS; 2) membrane transporters and ion channels involved in water as well as ion uptake; 3) transcription factors involved in controlling the transcription of stress related genes.

Different osmolytes have been reported to be associated with drought stress tolerance; enzymes involved in their metabolism represent potential candidates for generating drought stress tolerant transgenic plants (Sharma *et al.* 2011). Plants over-expressing proline biosynthesis genes have an increased tolerance to abiotic stress to some extent (Székely *et al.* 2008). Overproduction of proline in transgenic tobacco and other transgenic plants induces enhanced root biomass under limited water supply (Kishor *et al.* 1995). Transgenic soybean with the P5C reductase (*P5CR*) gene and the antisense construct from *Arabidopsis* prove that

proline enhances the survival during drought stress (De Ronde *et al.* 2000). It was found that two transgenic potato lines with enhanced expression of the yeast trehalose-6-phosphate synthase (*TPS1*) gene are more competent at maintaining water content and acceptable photosynthetic rate during drought in comparison with the wild type (Stiller *et al.* 2008). Glycine betaine biosynthetic genes in transgenic *Arabidopsis*, eucalypt, tobacco, rice, tomato, potato, and wheat plants proved to be highly effective in conferring stress tolerance (Goel *et al.* 2011). Several transgenic plants were produced using the genes involved in sorbitol biosynthesis. Transgenic tobacco plants expressing apple cDNA for sorbitol-6-phosphate-dehydrogenase accumulate a varying content of sorbitol. *Diospyros kaki* transformed with sorbitol-6-phosphate dehydrogenase accumulates sorbitol and shows a higher photosynthetic activity compared with control plants (Gao *et al.* 2001).

The increased activities of antioxidant enzymes obtained by the expression of foreign genes for enhancing abiotic stress tolerance have been reported in several plant species (Gupta *et al.* 1993, Allen *et al.* 1997, Payton *et al.* 1997, McKersie *et al.* 2000, Chen and Gallie 2005). Reports have also shown that *SOD* over-expression leads

to enhanced tolerance to drought stress (Bohnert and Sheveleva 1998). Alfalfa plants overexpressing the *Mn-SOD* from *Nicotiana plumbaginifolia* have been shown to have an improved yield and enhanced vigour after water deficit (McKersie *et al.* 1996). Over-expression of the *Populus* peroxisomal ascorbate peroxidase (*PpAPX*) gene in transgenic tobacco enhances its drought resistance (Li *et al.* 2009).

Expression of aquaporins was elevated in roots of rice plants grown in the presence of NH_4 , resulting in increased water permeability along with hydraulic conductivity (Ding *et al.* 2015). Over-expression of wheat *TaAQP7* confers drought stress tolerance to transgenic tobacco by enhancing water retention, reducing ROS accumulation by enhancing the enzymatic and non-enzymatic antioxidants, and reducing the membrane damage by decreasing lipid peroxidation (Zhou *et al.* 2012). Over-expression of the *Musa acuminata* aquaporin gene *PIP1;1* in *Arabidopsis* results in an increased primary root elongation along with root hair numbers and survival rates compared to the wild type under drought

stress (Xu *et al.* 2014).

Transcription activator and suppressor factors are key regulators involved in the induction of changes in the expression of stress related genes during the response to environmental stresses (Sakuma *et al.* 2006). Most of the transcription factors were identified and analysed in *Arabidopsis* using genome-wide microarrays to identify several potential target genes (Ravikumar *et al.* 2014). Over-expression of ABA stress and ripening induced protein 1 (*TaASR1*) in tobacco results in a decreased lipid peroxidation and membrane damage, as well as higher relative water content and antioxidant enzyme activities than in the wild type under drought stress (Hu *et al.* 2013). Ectopic expressions of *DREB* genes in *Arabidopsis*, wheat, barley, soybean, tomato, tobacco, strawberry, rice, oilseed rape, potato, *etc.*, enhance tolerance to several abiotic stresses (Hussain *et al.* 2011). Constitutive expression of *Arabidopsis DREB1A* and *ABF3* genes in transgenic rice increase tolerance to drought stress (Oh *et al.* 2005). MYB-type proteins are involved in diverse processes of plant growth,

Table 4. Expression of genes in different transgenic plants for enhanced drought tolerance.

Gene	Plant species	Function	Reference
<i>CjPIP2;1</i>	<i>Arabidopsis</i>	enhanced drought tolerance	Jang <i>et al.</i> 2007
<i>Dehydrin</i>	<i>Arabidopsis</i>	enhanced osmotic and salt stress tolerance	Hanin <i>et al.</i> 2011
<i>PIP2;2</i>	<i>Arabidopsis</i>	enhanced root water uptake capacity	Javot <i>et al.</i> 2003
<i>PgTIP1</i>	<i>Arabidopsis</i>	improved drought tolerance	Peng <i>et al.</i> 2007
<i>Rab 17</i>	<i>Arabidopsis</i>	enhanced osmotic stress tolerance	Figueras <i>et al.</i> 2004
<i>RcHSP17.8</i>	<i>Arabidopsis</i>	drought, salt, and osmotic stress tolerance	Jiang <i>et al.</i> 2009
<i>MIZ1</i>	<i>Arabidopsis</i>	dehydration avoidance under drought	Iwata <i>et al.</i> 2013
<i>CBL1</i>	<i>Arabidopsis</i>	enhanced drought tolerance	Cheong <i>et al.</i> 2003
<i>ABP9</i>	<i>Arabidopsis</i>	improved photosynthetic activity under drought stress	Zhang <i>et al.</i> 2008a
<i>OsWRKY45</i>	<i>Arabidopsis</i>	enhanced drought tolerance	Qiu <i>et al.</i> 2009
<i>AtHSP17.6A</i>	<i>Arabidopsis</i>	improved drought tolerance	Sun <i>et al.</i> 2001
<i>RWC3</i>	rice	maintenance of water potential and transcription during stress	Lian <i>et al.</i> 2004
<i>OsLEA3</i>	rice	enhanced dehydration tolerance	Zhang <i>et al.</i> 2009
<i>OsLEA3-1</i>	rice	improved drought resistance	Xiao <i>et al.</i> 2007
<i>HVA1</i>	rice	enhanced cell membrane stability	Babu <i>et al.</i> 2004
<i>PF00477</i>	rice	improved dehydration tolerance	Cheng <i>et al.</i> 2002
<i>sHSP17.7</i>	rice	enhanced drought tolerance	Sato <i>et al.</i> 2008
<i>OsiSAP8</i>	rice	drought and salt stress tolerance	Kanneganti <i>et al.</i> 2008
<i>OCP11</i>	rice	enhanced crop yield during drought	Huang <i>et al.</i> 2007
<i>HVA1</i>	rice	improved drought and salinity tolerance	Xu <i>et al.</i> 1996
<i>SodERF3</i>	tobacco	increased drought tolerance	Trujillo <i>et al.</i> 2009
<i>NtAQP1</i>	tobacco	increased root hydraulic conductance and drought tolerance	Siefritz <i>et al.</i> 2002
<i>LEA 4</i>	tobacco	improved membrane stability	Liu <i>et al.</i> 2009
<i>CaLEA6</i>	tobacco	protection of photosynthetic activity during drought stress	Jun <i>et al.</i> 2008
<i>DQ663481</i>	tobacco	improved cell membrane stability	Wang <i>et al.</i> 2006
<i>NtHSP70-1</i>	tobacco	regulated water flux and enhanced drought tolerance	Cho <i>et al.</i> 2006
<i>HVA1</i>	wheat	increased WUE and biomass	Sivamani <i>et al.</i> 2000
<i>LEA3-L2</i>	wheat	increased protein synthesis	NDong <i>et al.</i> 2002
<i>HVA1</i>	wheat	increased yield under drought stress	Bahieldin <i>et al.</i> 2005
<i>PIP</i>	soybean	enhanced drought tolerance	Porcel <i>et al.</i> 2006
<i>HVA1</i>	mulberry	enhanced drought and salt tolerance	Checker <i>et al.</i> 2012
<i>ME-leaN4</i>	lettuce	delayed wilting and improved drought tolerance	Park <i>et al.</i> 2005b
<i>ME-leaN4</i>	chinese cabbage	improved drought tolerance	Park <i>et al.</i> 2005a
<i>CaLEA</i>	chinese cabbage	enhanced dehydration tolerance	Park <i>et al.</i> 2003
<i>HVA1</i>	oat	delayed wilting under drought stress	Maqbool <i>et al.</i> 2002
<i>NPK1</i>	maize	protect photosynthetic activity during drought	Shou <i>et al.</i> 2004

development, and abiotic stress response. Transgenic *Arabidopsis* plants overexpressing the *TaMYB30-B* gene shows an altered expression of drought stress-responsive genes leading to improved drought stress tolerance during

the germination and the seedling stages (Zhang *et al.* 2012). A list of genes used for generating the transgenic plants with enhanced drought stress tolerance has been enclosed in Table 5.

Conclusions and future perspectives

Drought stress is one of the harsh environmental factors limiting the production of crop plants globally. The main challenge is to increase the production of plants under decreasing water potential, which can be achieved by understanding the changes in the osmotic potentials that in turn trigger an array of metabolic responses due to the increased expression of genes responsible for the protection of plants under various environmental conditions. The capability of drought stress tolerance in crop plants is more related to the adaptation of plants pertaining to changes in the environment and flexibility of their cellular metabolism, which is considered as key step in tolerance to environmental stresses. Changes in growth and development of plants are always related to changes in their physiological, biochemical, and molecular mechanisms (Fig. 1). For the past two decades, many drought stress inducible genes have been identified in crop plants after the genome sequencing of *Arabidopsis*. In spite of this progress, a critical gaps still remain in identifying mechanisms concerning phytohormone dependent and independent gene expression, and the role of enzymes in regulating toxicity of reactive nitrogen species and their metabolism along with identifying the rate limiting enzymes for the synthesis of osmo-protectants. Water deficit induces the expression of specific genes involved in adaptation of plants to stresses; the functions of some of them are not yet established. One way of understanding these functions is by identifying the key rate limiting steps that help in providing drought stress tolerance.

The ROS are important for regulating many cellular processes related to the response to changes in the environment. The ROS were initially recognized as toxic byproducts of an aerobic metabolism eliminated by the coordination of enzymatic and non-enzymatic antioxidants. Apart from their toxicity, the critical role of ROS includes their function as important signaling molecules in different processes involved in growth, development, responses to abiotic and biotic stress stimuli, and programmed cell death. Reactive oxygen species signaling is controlled by the balance of their production and scavenging similarly to calcium signaling which is controlled by storage and release. Still a few connections are missing, regarding the pathways that maintain optimum content of ROS in cells and the role of ROS in mediating the pathways oriented with plant protection mechanisms. Also, the role of ROS in cell to

cell communications and integration of networks related to the abiotic stress response and the possible interactions between ROS and reactive nitrogen species is still to be answered. Focus on these aspects will give comprehensive information on understanding drought tolerance mechanisms, and the application of these parameters on crop plants at a global level will lead to the production of crop plants with enhanced drought stress tolerance.




Fig. 1. Drought tolerance mechanisms in plants.

References

- Abe, H., Urao, T., Ito, T., Seki, M., Shinozaki, K., Yamaguchi-Shinozaki, K.: *Arabidopsis* AtMYC2 (bHLH) and AtMYB2 (MYB) functions as transcriptional activators in abscisic acid signaling. - *Plant Cell* **15**: 63-78, 2003.
- Adriano, S., Scopa, A., Nuzzaci, M., Vitti, A.: Ascorbate peroxidase and catalase activities and their genetic regulation in plants subjected to drought and salinity stresses. - *Int. J. mol. Sci.* **16**: 13561-13578, 2015.
- Ahmad, F., Rahmatullah, A.T., Maqsood, A.M., Tahir, M.A., Kanwal, S.: Effect of silicon application on wheat (*Triticum aestivum* L.) growth under water deficiency stress. - *Emir. J. Food Agr.* **19**: 1-7, 2007.
- Ahmad, P., Sarwat, M., Sharma, S.: Reactive oxygen species, antioxidants and signaling in plants. - *J. Plant Biol.* **51**: 167-173, 2008.
- Alexander, E., Frayssé, L., Sjövall-Larsen, S., Gustavsson, S., Fellert, M., Karlsson, M., Johanson, U., Kjellbom, P.: Whole gene family expression and drought stress regulation of aquaporins. - *Plant. mol. Biol.* **59**: 469-484, 2005.
- Allen, R.D., Webb, R.P., Schake, S.A.: Use of transgenic plants to study antioxidant defenses. - *Free Radical Biol. Med.* **23**: 473-479, 1997.
- Apel, H., Hirt, H.: Reactive oxygen species: metabolism, oxidative stress, and signal transduction. - *Annu. Rev. Plant Biol.* **55**: 373-399, 2004.
- Armin, S.M., Avat, S., Pessarakli, M.: Reactive oxygen species (ROS) generation and detoxifying in plants. - *J. Plant Nutr.* **37**: 1573-1585, 2014.
- Arora, A., Byrem, T.M., Nair, M.G., Strasburg, G.M.: Modulation of liposomal membrane fluidity by flavonoids and isoflavonoids. - *Arch. Biochem. Biophys.* **373**: 102-109, 2000.
- Asada, K.: The water-water cycle in chloroplasts: scavenging of active oxygens and dissipation of excess photons. - *Annu. Rev. Plant Physiol. Plant mol. Biol.* **50**: 601-639, 1999.
- Ashraf, M., Harris, P.J.C.: Photosynthesis under stressful environments: an overview. - *Photosynthetica* **51**: 163-190, 2013.
- Assefa, Y., Staggenborg, S.A., Prasad, V.P.V.: Grain sorghum water requirement and responses to drought stress: A review. - *Crop Manage. Online* **9**: doi:10.1094/CM-2010-1109-01-RV, 2010.
- Athar, H., Ashraf, M.: Photosynthesis under drought stress. - In: Pessarakli, M. (ed.) *Handbook of Photosynthesis*. 2nd Ed. Pp. 793-804. CRC Press, Taylor & Francis Group, New York 2005.
- Augé, R.M., Kubiková, E., Moore, J.L.: Foliar dehydration tolerance of mycorrhizal cowpea, soybean and bush bean. - *New Phytol.* **151**: 535-541, 2001.
- Babu, R.C., Zhang, J., Blum, A., Ho, T.H.D., Wu, R., Nguyen, H.T.: *HVA1*, a lea gene from barley confers dehydration tolerance in transgenic rice (*Oryza sativa* L.) via cell membrane protection. - *Plant Sci.* **166**: 855-862, 2004.
- Bahieldin, A., Mahfouz, H.T., Eissa, H.F., Saleh, O.M., Ramadan, A.M., Ahmed, I.A., Dyer, W.E., El-Itriby, H.A., Madkour, M.A.: Field evaluation of transgenic wheat plants stably expressing the *HVA1* gene for drought tolerance. - *Physiol. Plant.* **123**: 421-427, 2005.
- Bartels, D., Phillips, J., Chandler, J.: Desiccation tolerance: gene expression, pathways, and regulation of gene expression. - In: Jenks, M.A., Wood, A.J. (ed.): *Plant Desiccation Tolerance*. Pp. 115-137. Blackwell, Ames 2007.
- Barthakur, S., Babu, V., Bansal, K.C.: Overexpression of osmotin induces proline accumulation and confers tolerance to osmotic stress in transgenic tobacco. - *J. Plant Biochem. Biotechnol.* **10**: 31-37, 2001.
- Baum, J.A., Scandalios, J.G.: Developmental expression and intracellular location of superoxide dismutases in maize. - *Differentiation* **13**: 133-140, 1979.
- Berova, M., Zlatev, Z.: Influence of soil drought on growth and biomass partitioning in young bean (*Phaseolus vulgaris* L.) plants. - *Annu. Rep. Bean Improvement Coop.* **45**: 190-191, 2002.
- Blum, A.: Towards a conceptual ABA ideotype in plant breeding for water limited environments. - *Funct. Plant Biol.* **42**: 502-513, 2015.
- Bohnert, H.J., Sheveleva, E.: Plant stress adaptations, making metabolism move. - *Curr. Opin. Plant Biol.* **1**: 267-274, 1998.
- Boyer, J.S.: Leaf enlargement and metabolic rates in corn, soybean, and sunflower at various leaf water potentials. - *Plant Physiol.* **46**: 233-235, 1970.
- Boyer, J.S., James, R.A., Munns, R., Condon, T.G., Passioura, B.: Osmotic adjustment leads to anomalously low estimates of relative water content in wheat and barley. - *Funct. Plant Biol.* **35**: 1172-1182, 2008.
- Brossa, R., López-Carbonell, M., Jubany-Mario, T., Alegre, L.: Interplay between abscisic acid and jasmonic acid and its role in water-oxidative stress in wild-type, ABA-deficient, JA-deficient, and ascorbate-deficient *Arabidopsis* plants. - *J. Plant Growth Regul.* **30**: 322-333, 2011.
- Budak, H., Kantar, M., Kurtoglu, K.Y.: Drought tolerance in modern and wild wheat. - *Sci. World J.* **2013**: 548246-548262, 2013.
- Bunnag, S., Pongthai, P.: Selection of rice (*Oryza sativa* L.) cultivars tolerant to drought stress at the vegetative stage under field conditions. - *Amer. J. Plant Sci.* **4**: 1701-1708, 2013.
- Cannon, R.E., White, J.A., Scandalios, J.G.: Cloning of cDNA for maize superoxide dismutase (*SOD-2*). - *Proc. nat. Acad. Sci. USA* **84**: 179-183, 1987.
- Cha, J.Y., Kim, W.Y., Kang, S.B., Kim, J.I., Baek, D., Jung, I.J., Kim, M.R., Li, N., Kim, H.J., Nakajima, M., Asami, T., Sabir, J.S.M., Park, H.C., Lee, S.Y., Bohnert, H.J., Bressan, R.A., Pardo, J.M., Yun, D.J.: A novel thiol-reductase activity of *Arabidopsis* YUC6 confers drought tolerance independently of auxin biosynthesis. - *Nat. Commun.* **6**: 8041, 2015.
- Chastain, C.J., Heck, J.W., Colquhoun, T.A., Voge, D.G., Gu, X.Y.: Posttranslational regulation of pyruvate, orthophosphate dikinase in developing rice (*Oryza sativa*) seeds. - *Planta* **224**: 924-934, 2006.
- Chaumont, F., Tyerman, S.D.: Aquaporins: highly regulated channels controlling plant water relations. - *Plant Physiol.* **164**: 1600-1618, 2014.
- Chaves, M.M., Flexas, J., Pinheiro, C.: Photosynthesis under drought and salt stress: regulation mechanisms from whole plant to cell. - *Ann. Bot.* **103**: 551-560, 2009.
- Checker, V.G., Chhibbar, A.K., Khurana, P.: Stress-inducible expression of barley *Hva1* gene in transgenic mulberry displays enhanced tolerance against drought, salinity and cold stress. - *Transgenic Res.* **21**: 939-957, 2012.
- Chen, T.H.H., Murata, N.: Glycine betaine: an effective protectant against abiotic stress in plants. - *Trends Plant Sci.* **13**: 499-505, 2008.
- Chen, Z., Gallie, D.R.: Increasing tolerance to ozone by elevating foliar ascorbic acid confers greater protection against ozone than increasing avoidance. - *Plant Physiol.* **138**: 1673-1689,

- 2005.
- Cheng, Z.Q., Targolli, J., Huang, X.Q., Wu, R.: Wheat LEA genes, *PMA80* and *PMA1959*, enhance dehydration tolerance of transgenic rice (*Oryza sativa* L.). - Mol. Breed. **10**: 71-82, 2002.
- Cheong, Y.H., Kim, K., Pandey, G.K., Gupta, R., Grant, J.J., Luan, S.: CBL1, a calcium sensor that differentially regulates salt, drought, and cold responses in *Arabidopsis*. - Plant Cell **15**: 1833-1845, 2003.
- Cho, E.K., Hong, C.B.: Over-expression of tobacco *NtHSP70-1* contributes to drought-stress tolerance in plants. - Plant Cell Rep. **25**: 349-358, 2006.
- Cobbett, C.S.: Phytochelators and their roles in heavy metal detoxification. - Plant Physiol. **123**: 825-832, 2000.
- Collins, A.: Carotenoids and genomic stability. - Mutat. Res. **475**: 21-28, 2001.
- Cortina, C., Culiáñez-Macià, F.A.: Tomato abiotic stress enhanced tolerance by trehalose biosynthesis. - Plant Sci. **169**: 75-82, 2005.
- Dat, J., Vandenabeele, S., Vranova, E., Van Montagu, M., Inze, D., Van Breusegem, F.: Dual action of the active oxygen species during plant stress responses. - Cell. Mol. Life Sci. **57**: 779-795, 2000.
- De Ronde, J.A., Spreeth, M.H., Cress, W.A.: Effect of antisense L- Δ^1 -pyrroline-5-carboxylate reductase transgenic soybean plants subjected to osmotic and drought stress. - Plant Growth Regul. **32**: 13-26, 2000.
- Ding, L., Gao, C., Li, Y., Li, Y., Zhu, Y., Xu, G., Shen, Q., Kaldenhoff, R., Kai, L., Guo, S.: The enhanced drought tolerance of rice plants under ammonium is related to aquaporin (AQP). - Plant Sci. **234**: 14-21, 2015.
- Dixon, D.P., Laphorn, A., Edwards, R.: Plant glutathione transferases. - Genome Biol. **3**: 3004.1-3004.10, 2002.
- Dong, C., Zheng, X., Li, G., Zhu, H., Zhou, M., Hu, Z.: Molecular cloning and expression of two cytosolic copper-zinc superoxide dismutases genes from *Nelumbo nucifera*. - Appl. Biochem. Biotechnol. **163**: 679-691, 2011.
- Doubnerová, V., Ryšlavá, H.: What can enzymes of C4 photosynthesis do for C3 plants under stress? - Plant Sci. **180**: 575-583, 2011.
- Du, Y.C., Nose, A., Wasano, K., Uchida, Y.: Responses to water stress of enzyme activities and metabolite levels in relation to sucrose and starch synthesis, the Calvin cycle and the C4 pathway in sugarcane (*Saccharum* sp.) leaves. - Aust. J. Plant Physiol. **25**: 253-260, 1998.
- Dubois, M., Skirycz, A., Claeys, H., Maleux, K., Dhondt, S., Van den Bossche, R., De Milde, L., Yoshizumi, T., Matsui, M., Inzé, D.: Ethylene response Factor 6 acts as central regulator of leaf growth under water-limiting conditions in *Arabidopsis*. - Plant Physiol. **162**: 319-332, 2013.
- Eltayeb, A.E., Kawano, N., Badawi, G.H., Kaminaka, H., Sanekata, T., Shibahara, T., Inanaga, S., Tanaka, K.: Overexpression of monodehydroascorbate reductase in transgenic tobacco confers enhanced tolerance to ozone, salt and polyethylene glycol stresses. - Planta **225**: 1255-1264, 2007.
- Fariduddin, Q., Varshney, P., Yusuf, M., Ali, A., Ahmad, A.: Dissecting the role of glycine betaine in plants under abiotic stress. - Plant Stress **7**: 8-18, 2013.
- Farooq, M., Wahid, A., Kobayashi, N., Fujita, D., Basra, S.M.A.: Plant drought stress: effects, mechanisms and management. - Agron. Sustainable Dev. **29**: 185-212, 2009.
- Figueras, M., Pujal, J., Saleh, A., Save, R., Pagès, M., Goday, A.: Maize *Rab17* overexpression in *Arabidopsis* plants promotes osmotic stress tolerance. - Ann. appl. Biol. **144**: 251-257, 2004.
- Flexas, J., Bota, J., Loreto, F., Cornic, G., Sharkey, T.D.: Diffusive and metabolic limitations to photosynthesis under drought and salinity in C3 plants. - Plant Biol. **6**: 269-279, 2004.
- Foyer, C.H., Noctor, G.: Oxygen processing in photosynthesis: regulation and signalling. - New Phytol. **146**: 359-388, 2000.
- Galau, G.A., Wang, H.Y.C., Hughes, D.W.: Cotton *Lea5* and *Lea14* encode atypical late embryogenesis abundant proteins. - Plant Physiol. **101**: 695-696, 1993.
- Galmes, J., Pou, A., Alsina, M.M., Tomàs, M., Medrano, H., Flexas, J.: Aquaporin expression in response to different water stress intensities and recovery in Richter-110 (*Vitis* sp): relationship with ecophysiological status. - Planta **226**: 671-681, 2007.
- Gao, M., Tao, R., Miura, K., Dandekar, A.M., Sugiura, A.: Transformation of Japanese persimmon (*Diospyros kaki* Thunb.) with apple cDNA encoding NADP-dependent sorbitol-6-phosphate dehydrogenase. - Plant Sci. **160**: 837-845, 2001.
- Ghotbi-Ravandi, A.A., Shahbazi, M., Shariati, M., Mulo, P.: Effects of mild and severe drought stress on photosynthetic efficiency in tolerant and susceptible barley (*Hordeum vulgare* L.) genotypes. - J. Agron. Crop Sci. **200**: 403-415, 2014.
- Giovannoni, J.J.: Completing a pathway to plant vitamin C synthesis. - Proc. nat. Acad. Sci. USA **104**: 9109-9110, 2007.
- Giuliani, S., Clarke, J., Kreps, J., Sanguinetti, M.C., Salvi, S., Landi, P., Zhu, T., Tuberosa, R.: Microarray analysis of backcrossed-derived lines differing for root-ABA1, a major QTL controlling root characteristics and ABA concentration in maize. - In: Tuberosa, R., Phillips, R.L., Gale, M. (ed.): Proceedings of the International Congress in the Wake of the Double Helix: From the Green Revolution to the Gene Revolution. Pp. 463-490. Avenue Media, Bologna 2005.
- Goddijn, O.J., Verwoerd, T.C., Voogd, E., Krutwagen, R.W., De Graaf, P.T., Van Dun, K., Poels, J., Ponstein, A.S., Damm, B., Pen, J.: Inhibition of trehalase activity enhances trehalose accumulation in transgenic plants. - Plant Physiol. **113**: 181-190, 1997.
- Goel, D., Singh, A.K., Yadav, V., Babbar, S.B., Bansa, K.C.: Overexpression of osmotin gene confers tolerance to salt and drought stresses in transgenic tomato (*Solanum lycopersicum* L.). - Protoplasma **245**: 133-141, 2010.
- Goel, D., Singh, A.K., Yadav, V., Babbar, S.B., Murata, N., Bansal, K.C.: Transformation of tomato with a bacterial *codA* gene enhances tolerance to salt and water stresses. - J. Plant Physiol. **168**: 1286-1294, 2011.
- Gomes, D., Agasse, A., Thiébaud, P., Delrot, S., Gerós, H., Chaumont, F.: Aquaporins are multifunctional water and solute transporters highly divergent in living organisms. - Biochim. biophys. Acta **1788**: 1213-1228, 2009.
- Grace, S.G., Logan, B.A.: Energy dissipation and radical scavenging by the plant phenylpropanoid pathway. - Phil. Trans. roy. Soc. London B **355**: 1499-1510, 2000.
- Grelet, J., Benamar, A., Teyssier, E., Avelange-Macherel, M.H., Grunwald, D., Macherel, D.: Identification in pea seed mitochondria of a late-embryogenesis abundant protein able to protect enzymes from drying. - Plant Physiol. **137**: 157-167, 2005.
- Grillo, S., Leone, A., Xu, Y., Tucci, M., Francione, R., Hasegawa, P.M., Monti, L., Bressan, R.A.: Control of osmotin gene expression by ABA and osmotic stress in vegetative tissues of wild-type and ABA-deficient mutants of tomato. - Physiol. Plant. **93**: 498-504, 1995.

- Guler, N.S., Saglam, A., Demiralay, M., Kadioglu, A.: Apoplastic and symplastic solute concentrations contribute to osmotic adjustment in bean genotypes during drought stress. - *Turk. J. Biol.* **36**: 151-160, 2012.
- Guo, P.G., Baum, M., Grando, S., Ceccarelli, S., Bai, G.H., Li, R.H., Von Korff, M., Varshney, R.K., Graner, A., Valkoun, J.: Differentially expressed genes between drought-tolerant and drought-sensitive barley genotypes in response to drought stress during the reproductive stage. - *J. exp. Bot.* **60**: 3531-3544, 2009.
- Gupta, N., Thind, S.: Improving photosynthetic performance of bread wheat under field drought stress by foliar applied glycine betaine. - *J. Agr. Sci. Technol.* **17**: 75-86, 2015.
- Gupta, S.A., Heinen, J.L., Holaday, A.S., Burke, J.J., Allen, R.D.: Increased resistance in transgenic plants that overexpress chloroplastic Cu/Zn superoxide dismutase. - *Proc. nat. Acad. Sci. USA* **90**: 1629-1633, 1993.
- Halliwel, B.: Oxidative stress and neurodegeneration: where are we now? - *J. Neurochem.* **97**: 1634-1658, 2006.
- Hancock, J., Desikan, R., Harrison, J., Bright, J., Hooley, R., Neill, S.: Doing the unexpected: proteins involved in hydrogen peroxide perception. - *J. exp. Bot.* **57**: 1711-1718, 2006.
- Hanin, M., Brini, F., Ebel, C., Toda, Y., Takeda, S., Masmoudi, K.: Plant dehydrins and stress tolerance: versatile proteins for complex mechanisms. - *Plant Signal Behav.* **6**: 1503-1509, 2011.
- Hare, P.D., Cress, W.A.: Metabolic implications of stress induced proline accumulation in plants. - *Plant Growth Regul.* **21**: 79-102, 1997.
- Hernández, I., Cela, J., Alegre, L., Munne-Bosch, S.: Antioxidant defenses against drought stress. - In: Aroca, R. (ed.): *Plant Responses to Drought Stress: from Morphological to Molecular Features*. Vol. 1. Pp. 231-258. Springer-Verlag, Berlin - Heidelberg 2012.
- Hu, W., Huang, C., Deng, X., Zhou, S., Chen, L., Li, Y., Wang, C., Ma, Z., Yuan, Q., Wang, Y., Cai, R., Liang, X., Yang, G., He, G.: *TaASR1*, a transcription factor gene in wheat, confers drought stress tolerance in transgenic tobacco. - *Plant Cell Environ.* **36**: 1449-64, 2013.
- Huang, Y., Xiao, B., Xiong, L.: Characterization of a stress responsive proteinase inhibitor gene with positive effect in improving drought resistance in rice. - *Planta* **226**: 73-85, 2007.
- Hussain, S.S., Kayani, M.A., Amjad, M.: Transcription factors as tools to engineer enhanced drought stress tolerance in plants. - *Biotechnol. Progr.* **27**: 297-306, 2011.
- Igamberdiev, A.U., Seregelyes, C., Manach, N., Hill, R.D.: NADH-dependent metabolism of nitric oxide in alfalfa root cultures expressing barley haemoglobin. - *Planta* **219**: 95-102, 2004.
- Iwata, S., Miyazawa, Y., Fujii, N., Takahashi, H.: MIZ1-regulated hydrotropism functions in the growth and survival of *Arabidopsis thaliana* under natural conditions. - *Ann. Bot.* **112**: 103-114, 2013.
- Jaleel, C.A., Manivannan, P., Wahid, A., Farooq, M., Somasundaram, R., Panneerselvam, R.: Drought stress in plants: a review on morphological characteristics and pigments composition. - *Int. J. agr. Biol.* **11**: 100-105, 2009.
- Jang, I.C., Oh, S.J., Seo, J.S., Choi, W.B., Song, S.I., Kim, C.H., Kim, Y.S., Seo, H.S., Choi, Y.D., Nahm, B.H., Kim, J.K.: Expression of a bifunctional fusion of the *Escherichia coli* genes for trehalose-6-phosphate synthase and trehalose-6-phosphate phosphatase in transgenic rice plants increases trehalose accumulation and abiotic stress tolerance without stunting growth. - *Plant Physiol.* **131**: 516-524, 2003.
- Jang, J.Y., Rhee, J.Y., Kim, D.G., Chung, G.C., Lee, J.H., Kang, H.: Ectopic expression of a foreign aquaporin disrupts the natural expression patterns of endogenous aquaporin genes and alters plant responses to different stress conditions. - *Plant Cell Physiol.* **48**: 1331-1339, 2007.
- Jaspers, P., Kangasjärvi, J.: Reactive oxygen species in abiotic stress signaling. - *Physiol. Plant.* **138**: 405-413, 2010.
- Javot, H., Laugeat, V., Santoni, V., Martin-Laurent, F., Güçlü, J., Vinh, J., Heyes, J., Franck, K.I., Schäffner, A.R., Bouchez, D., Maurel, C.: Role of a single aquaporin isoform in root water uptake. - *Plant Cell* **15**: 509-522, 2003.
- Jiang, C., Xu, J., Zhang, H., Zhang, X., Shi, J., Li, M., Ming, F.: A cytosolic class I small heat shock protein, RcHSP17.8, of *Rosa chinensis* confers resistance to a variety of stresses to *Escherichia coli*, yeast and *Arabidopsis thaliana*. - *Plant Cell Environ.* **32**: 1046-1059, 2009.
- Jun, S.S., Choi, H.J., Lee, H.Y., Hong, Y.N.: Differential protection of photosynthetic capacity in trehalose-and lea protein-producing transgenic plants under abiotic stresses. - *J. Plant Biol.* **51**: 327-336, 2008.
- Kanematsu, S., Asada, K.: Superoxide dismutase. - In: Fukui, T., Soda, K. (ed.): *Molecular Aspects of Enzyme Catalysis*. Pp. 191-209. Wiley, Weinheim 1994.
- Kannan, N.D., Kulandaivelu, G.: Drought induced changes in physiological, biochemical and phytochemical properties of *Withania somnifera* Dun. - *J. med. Plants Res.* **5**: 3929-3935, 2011.
- Kanneganti, V., Gupta, A.: Overexpression of *OsiSAP8*, a member of stress associated protein (*SAP*) gene family of rice confers tolerance to salt, drought and cold stress in transgenic tobacco and rice. - *Plant mol. Biol.* **66**: 445-462, 2008.
- Karuppanandian, T., Wang, H.W., Prabakaran, N., Jeyalakshmi, K., Kwon, M., Manoharan, K., Kim, W.: 2,4-dichlorophenoxyacetic acid-induced leaf senescence in mung bean (*Vigna radiata* L. Wilczek) and senescence inhibition by co-treatment with silver nanoparticles. - *Plant Physiol. Biochem.* **49**: 168-177, 2011.
- Kawamitsu, Y., Driscoll, T., Boyer, J.S.: Photosynthesis during desiccation in an intertidal alga and a land plant. - *Plant Cell Physiol.* **41**: 344-353, 2000.
- Keles, Y., Unyayar, S.: Responses of antioxidant defense system of *Helianthus annuus* to abscisic acid under drought and waterlogging. - *Acta Physiol. Plant.* **26**: 149-156, 2004.
- Kishor, K.P.B., Hong, Z., Miao, G.H., Hu, C.A.A., Verma, D.P.S.: Overexpression of [delta]-pyrroline-5-carboxylate synthetase increase proline production and confers osmotolerance in transgenic plants. - *Plant Physiol.* **108**: 1387-1394, 1995.
- Kliebenstein, D.J., Dietrich, R.A., Martin, A.C., Last, R.L., Dangel, J.L.: LSD1 regulates salicylic acid induction of copper zinc superoxide dismutase in *Arabidopsis thaliana*. - *Mol. Plant Microbe Interact.* **12**: 1022-1026, 1999.
- Koffler, B.E., Luschn-Ebengreuth, N., Stabenheiner, E., Müller, M., Zechmann, B.: Compartment specific response of antioxidants to drought stress in *Arabidopsis*. - *Plant Sci.* **227**: 133-144, 2014.
- Kopriva, S., Rennenberg, H.: Control of sulphate assimilation and glutathione synthesis: interaction with N and C metabolism. - *J. exp. Bot.* **55**: 1831-1842, 2004.
- Krouma, A.: Plant water relations and photosynthetic activity in three Tunisian chickpea (*Cicer arietinum* L.) genotypes subjected to drought. - *Turk. J. Agr. Forest.* **34**: 257-264, 2010.
- Kumar, R.R., Sharma, S.K., Goswami, S., Singh, K., Gadpayle,

- K.A., Singh, G.P., Pathak, H., Rai, R.D.: Transcript profiling and biochemical characterization of mitochondrial superoxide dismutase (mtSOD) in wheat (*Triticum aestivum*) under different exogenous stresses. - Aust. J. Crop Sci. **7**: 414-424, 2013.
- Lambert, J.D., Elias, R.J.: The antioxidant and pro-oxidant activities of green tea polyphenols: a role in cancer prevention. - Arch. Biochem. Biophys. **501**: 65-67, 2010.
- Lawlor, D.W.: Genetic engineering to improve plant performance under drought: physiological evaluation of achievements, limitations, and possibilities. - J. exp. Bot. **64**: 83-108, 2013.
- Li, Y.-J., Hai, R.-L., Du, X.-H., Jiang, X.-N., Lu, H.: Overexpression of a *Populus* peroxisomal ascorbate peroxidase (*PpAPX*) gene in tobacco plants enhances stress tolerance. - Plant Breed. **128**: 404-410, 2009.
- Lian, H.L., Yu, X., Ye, Q., Ding, X., Kitagawa, Y., Kwak, S.S., Su, W.A., Tang, Z.C.: The role of aquaporin RWC3 in drought avoidance in rice. - Plant Cell Physiol. **45**: 481-489, 2004.
- Liu, X., Wang, Z., Wang, L., Wu, R., Phillips, J., Deng, X.: *LEA 4* group genes from the resurrection plant *Boea hygrometrica* confer dehydration tolerance in transgenic tobacco. - Plant Sci. **176**: 90-98, 2009.
- Livingston, D.P., Hinch, D.K., Heyer, A.G.: Fructan and its relationship to abiotic stress tolerance in plants. - Cell. mol. Life Sci. **66**: 2007-2023, 2009.
- Lutts, S., Almansouri, M., Kinet, J.M.: Salinity and water stress have contrasting effects on the relationship between growth and cell viability during and after stress exposure in durum wheat. - Plant Sci. **167**: 9-18, 2004.
- Maqbool, S.B., Zhong, H., El-Maghraby, Y., Ahmad, A., Chai, B., Wang, W., Sabzikar, R., Sticklen, M.B.: Competence of oat (*Avena sativa* L.) shoot apical meristems for integrative transformation, inherited expression, and osmotic tolerance of transgenic lines containing hva1. - Theor. appl. Genet. **105**: 201-208, 2002.
- Maroco, J., Rodrigues, M.L., Lopes, C., Chaves, M.M.: Limitations to leaf photosynthesis in grapevine under drought-metabolic and modelling approaches. - Funct. Plant Biol. **29**: 451-459, 2002.
- Martinez, J.P., Silva, H., Ledent, J.F., Pinto, M.: Effect of drought stress on the osmotic adjustment, cell wall elasticity and cell volume of six cultivars of common beans (*Phaseolus vulgaris* L.). - Europ. J. Agron. **26**: 30-38, 2007.
- Maurel, C., Verdoucq, L., Luu, D.T., Santoni, V.: Plant aquaporins: membrane channels with multiple integrated functions. - Annu. Rev. Plant Biol. **59**: 595-624, 2008.
- McKersie, B.D., Bowley, S.R., Harjanto, E., Leprince, O.: Water-deficit tolerance and field performance of transgenic alfalfa overexpressing superoxide dismutase. - Plant Physiol. **111**: 1177-1181, 1996.
- McKersie, B.D., Murnaghan, J., Jones, K.S., Bowley, S.R.: Iron-superoxide dismutase expression in transgenic alfalfa increases winter survival without a detectable increase in photosynthetic oxidative stress tolerance. - Plant Physiol. **122**: 1427-1437, 2000.
- Merewitz, E.B., Gianfagna, T., Huang, B.: Photosynthesis, water use, and root viability under water stress as affected by expression of *SAG12*-ipt controlling cytokinin synthesis in *Agrostis stolonifera*. - J. exp. Bot. **62**: 383-395, 2011.
- Misra, A.N., Misra, M., Singh, R.: Chlorophyll fluorescence in plant biology. - In: Misra, A.N. (ed.): Biophysics. Pp. 171-192. InTech 2012.
- Mittler, R., Blumwald, E.: The roles of ROS and ABA in systemic acquired acclimation. - Plant Cell **27**: 64-70, 2015.
- Miyake, C., Cao, W.H., Asada, K.: Purification and molecular properties of thylakoid-bound ascorbate peroxidase from spinach chloroplasts. - Plant Cell Physiol. **34**: 881-889, 1993.
- Moloudi, F., Navabpour, S., Soltanloo, H., Ramazanpour, S.S., Sadeghipour, H.: Catalase and metallothionein genes expression analysis in wheat cultivars under drought stress condition. - J. Plant mol. Breed. **1**: 54-68, 2013.
- Moons, A., De Keyser, A., Van Montagu, M.: A group 3 *LEA* cDNA of rice, responsive to abscisic acid, but not to jasmonic acid, shows variety-specific differences in salt stress response. - Gene **191**: 197-204, 1997.
- Most, P., Jutta, P.: Possible roles of plant sulfurtransferases in detoxification of cyanide, reactive oxygen species, selected heavy metals and arsenate. - Molecules **20**: 1410-1423, 2015.
- Mumm, P., Wolf, T., Fromm, J., Rob, M., Roelfsema, G., Marten, I.: Cell type-specific regulation of ion channels within the maize stomatal complex. - Plant Cell Physiol. **52**: 1365-1375, 2011.
- Nakabayashi, R., Mori, T., Saito, K.: Alternation of flavonoid accumulation under drought stress in *Arabidopsis thaliana*. - Plant Signal. Behav. **9**: e29518, 2014.
- Naya, L., Ladrera, R., Ramos, J., González, E.M., Arrese-Igor, C., Minchin, F.R., Becana, M.: The response of carbon metabolism and antioxidant defenses of alfalfa nodules to drought stress and to the subsequent recovery of plants. - Plant Physiol. **144**: 1104-1114, 2007.
- NDong, C., Danyluk, J., Wilson, K.E., Pocock, T., Huner, N.P. A., Sarhan, F.: Cold-regulated cereal chloroplast late embryogenesis abundant-like proteins. Molecular characterization and functional analyses. - Plant Physiol. **129**: 1368-1381, 2002.
- Nikinmaa, E., Hölttä, T., Hari, P., Kolari, P., Mäkelä, A., Sevanto, S., Vesala, T.: Assimilate transport in phloem sets conditions for leaf gas exchange. - Plant, Cell Environ. **36**: 655-669, 2013.
- Oh, S.J., Song, S.I., Kim, Y.S., Jang, H.J., Kim, M., Kim, Y.K.: *Arabidopsis CBF3/DREB1A* and *ABF3* in transgenic rice increased tolerance to abiotic stress without stunting growth. - Plant Physiol. **138**: 341-351, 2005.
- Olien, C.R., Clark, J.L.: Freeze-induced changes in carbohydrates associated with hardiness of barley and rye. - Crop Sci. **35**: 496-502, 1995.
- Osakabe, Y., Osakabe, K., Shinozaki, K., Tran, L.S.P.: Response of plants to water stress. - Front. Plant Sci. **5**: 1-8, 2014.
- Padmavathi, T.A.V., Rao, D.M.: Differential accumulation of osmolytes in 4 cultivars of peanut (*Arachis hypogaea* L.) under drought stress. - J. Crop Sci. Biotechnol. **16**: 151-159, 2013.
- Pang, C.H., Wang, B.S.: Role of ascorbate peroxidase and glutathione reductase in ascorbate-glutathione cycle and stress tolerance in plants. - In: Anjum, A.N., Chan, M.T., Umar, S. (ed.): Ascorbate-Glutathione Pathway and Stress Tolerance in Plants. Pp. 99-113. Springer, Berlin 2010.
- Park, B.J., Liu, Z., Kanno, A., Kamey, T.: Genetic improvement of Chinese cabbage for salt and drought tolerance by constitutive expression of a *B. napus LEA* gene. - Plant Sci. **169**: 553-558, 2005a.
- Park, B.J., Liu, Z., Kanno, A., Kamey, T.: Increased tolerance to salt and water-deficit stress in transgenic lettuce (*Lactuca sativa* L.) by constitutive expression of *LEA*. - Plant Growth Regul. **45**: 165-171, 2005b.
- Park, S., Li, J., Pittman, J.K., Berkowitz, G.A., Yang, H., Undurraga, S., Morris, J., Hirschi, K.D., Gaxiola, R.A.: Up-regulation of a H⁺-pyrophosphatase (H⁺-PPase) as a strategy to engineer drought-resistant crop plants. - Proc. nat. Acad.

- Sci. USA. **102**: 18830-18835, 2005.
- Park, S.H., Jun, S.S., An, G., Hong, Y.N., Park, M.C.: A comparative study on the protective role of trehalose and LEA proteins against abiotic stresses in transgenic Chinese cabbage (*Brassica campestris*) overexpressing *CaLEA* or *OtsA*. - J. Plant Biol. **46**: 277-286, 2003.
- Parkhi, V., Kumar, V., Sunilkumar, G., Campbell, L.M., Singh, N.K., Rathore, K.S.: Expression of apoplastically secreted tobacco osmotin in cotton confers drought tolerance. - Mol. Breed. **23**: 625-639, 2009.
- Payton, P., Allen, R.D., Trolinder, N., Holaday, A.S.: Overexpression of chloroplast-targeted Mn superoxide dismutase in cotton (*Gossypium hirsutum* L., cv. Coker 312) does not alter the reduction of photosynthesis after short exposures to low temperature and high light intensity. - Photosynth. Res. **52**: 233-244, 1997.
- Pei, E.Q., Dan, Y.S., Liu, P.Q., Song, Y.C., Wang, T.Y., Li, Y.: Effect of drought stress on osmotic adjustment in different maize inbred lines at seedling stage. - J. Plant Genet. Resour. **11**: 40-45, 2010.
- Peña-Valdivia, C.B., Sánchez-Urdaneta, A.B., Meza, R.J., Juárez, M.J., García-Nava, R., Celis Velázquez, R.: Anatomical root variations in response to water deficit: wild and domesticated common bean (*Phaseolus vulgaris* L.). - Biol. Res. **43**: 417-427, 2010.
- Peng, Y., Lin, W., Cai, W., Arora, R.: Overexpression of a *Panax ginseng* tonoplast aquaporin alters salt tolerance, drought tolerance and cold acclimation ability in transgenic *Arabidopsis* plants. - Planta **226**: 729-40, 2007.
- Porcel, R., Aroca, R., Azcn, R., Ruiz-Lozano, J.M.: PIP aquaporin gene expression in arbuscular mycorrhizal *Glycine max* and *Lactuca sativa* plants in relation to drought stress tolerance. - Plant mol. Biol. **60**: 389-404, 2006.
- Praba, M.L., Cairns, J.E., Babu, R.C., Lafitte, H.R.: Identification of physiological traits underlying cultivar differences in drought tolerance in rice and wheat. - J. Agron. Crop. Sci. **195**: 30-46, 2009.
- Qiu, Y., Yu, D.: Over-expression of the stress-induced *OsWRKY45* enhances disease resistance and drought tolerance in *Arabidopsis*. - Environ. exp. Bot. **65**: 35-47, 2009.
- Raghavendra, A.S., Gonugunta, V.K., Christmann, A., Grill, E.: ABA perception and signalling. - Trends Plant Sci. **15**: 395-401, 2010.
- Raghothama, K.G., Liu, D., Nelson, D.E., Hasegawa, P.M., Bressan, R.A.: Analysis of an osmotically regulated pathogenesis-related osmotin gene promoter. - Plant mol. Biol. **23**: 1117-1128, 1993.
- Rahdari, P., Hoseini, S.M.: Drought Stress: a review. - Int. J. Agron. Plant Prod. **3**: 443-446, 2012.
- Ramos, M.L.G., Gordon, A.J., Minchin, F.R., Sprent, J.J., Parsons, R.: Effect of water stress on nodule physiology and biochemistry of a drought tolerant cultivar of common bean (*Phaseolus vulgaris* L.). - Ann. Bot. **83**: 57-63, 1999.
- Rao, D.E., Divya, K., Ramana, G.V., Chaitanya, K.V.: Biochemical and molecular mechanisms that decipher drought stress tolerance in plants. - In: Gupta, R.K., Akhtar, N., Vyas, D. (ed.): Biotechnology: an Overview. Pp. 387-398. Daya Publishing House, New Delhi 2015.
- Rauf, S., Sadaqat, H.A.: Effect of osmotic adjustment on root length and dry matter partitioning in sunflower (*Helianthus annuus* L.) under drought stress. - Acta. agr. scand. **58**: 252-260, 2008.
- Ravikumar, G., Manimaran, P., Voleti, S.R., Subrahmanyam, D., Sundaram, R.M., Bansal, K.C., Viraktamath, B.C., Balachandran, S.M.: Stress-inducible expression of *AtDREB1A* transcription factor greatly improves drought stress tolerance in transgenic indica rice. - Transgenic Res. **23**: 421-439, 2014.
- Raynal, M., Gaubier, P., Grellet, F., Delseny, M.: Nucleotide sequence of a radish cDNA clone coding for a late embryogenesis abundant (LEA) protein. - Nucl. Acids Res. **18**: 6132, 1990.
- Reddie, K.G., Carroll, K.S.: Expanding the functional diversity of proteins through cysteine oxidation. - Curr. Opin. Chem. Biol. **12**: 746-754, 2008.
- Reddy, A.R., Chaitanya, K.V., Vivekanandan, M.: Drought-induced responses of photosynthesis and antioxidant metabolism in higher plants. - J. Plant Physiol. **161**: 1189-1202, 2004.
- Rhodes, D., Hanson, A.D.: Quaternary ammonium and tertiary sulfonium compounds in higher plants. - Annu. Rev. Plant Physiol. **44**: 357-384, 1993.
- Rosa, S.B., Caverzan, A., Teixeira, F.K., Lazzarotto, F., Silveira, J.A., Ferreira-Silva, S.L., Abreu-Neto, J., Margis, R., Margis-Pinheiro, M.: Cytosolic APX knockdown indicates an ambiguous redox responses in rice. - Phytochemistry **71**: 548-558, 2010.
- Rouhier, N., Couturier, J., Jacquot, J.P.: Genome-wide analysis of plant glutaredoxin systems. - J. exp. Bot. **57**: 1685-1696, 2006.
- Saeidnejad, A.H., Rajaei, P.: Antioxidative responses to drought and salinity stress in plants, a comprehensive review. - Int. J. Life Sci. **9**: 1-8, 2015.
- Sage, R.F., Sage, T.L., Kocacinar, F.: Photorespiration and the evolution of C₄ photosynthesis. - Annu. Rev. Plant Biol. **63**: 19-47, 2012.
- Sakamoto, H., Maruyama, K., Sakuma, Y., Meshi, T., Iwabuchi, M., Shinozaki, K., Yamaguchi-Shinozaki, K.: *Arabidopsis* Cys2His2-type zinc-finger proteins function as transcription repressors under drought, cold and high-salinity stress conditions. - Plant Physiol. **136**: 2734-2746, 2004.
- Sakuma, Y., Maruyama, K., Osakabe, Y., Qin, F., Seki, M., Shinozaki, K., Yamaguchi-Shinozaki, K.: Functional analysis of an *Arabidopsis* transcription factor, DREB2A, involved in drought-responsive gene expression. - Plant Cell **18**: 1292-1309, 2006.
- Sakuraba, Y., Kim, Y.S., Han, S.H., Lee, B.D., Paek, N.C.: The *Arabidopsis* transcription factor NAC016 promotes drought stress responses by repressing AREB1 transcription through a trifurcate feed-forward regulatory loop involving NAP. - Plant Cell **27**: 1771-1787, 2015.
- Samarah, N.H., Alqudah, A.M., Amayreh, J.A., McAndrews, G.M.: The effect of late-terminal drought stress on yield components of four barley cultivars. - J. Agron. Crop. Sci. **195**: 427-441, 2009.
- Sato, Y., Yokoya, S.: Enhanced tolerance to drought stress in transgenic rice plants overexpressing a small heat-shock protein, sHSP17.7. - Plant Cell Rep. **27**: 329-334, 2008.
- Schippers, J.H.M., Nguyen, H.M., Lu, D., Schmidt, R., Mueller-Roeber, B.: ROS homeostasis during development: an evolutionary conserved strategy. - Cell Mol. Life Sci. **69**: 3245-3257, 2012.
- Sharma, P., Jha, A.B., Dubey, R.S., Pessarakli, M.: Reactive oxygen species generation, hazards, and defense mechanisms in plants under environmental (abiotic and biotic) stress conditions. - In: Pessarakli, M. (ed.): Handbook of Plant and Crop Physiology. 3rd Ed. Pp. 509-548. CRC Press, Taylor & Francis Group, New York 2014.
- Sharma, S., Villamor, J.G., Verslues, P.E.: Essential role of

- tissue-specific proline synthesis and catabolism in growth and redox balance at low water potential. - *Plant Physiol.* **157**: 292-304, 2011.
- Shih, M.D., Lin, S.C., Hsieh, J.S., Tsou, C.H., Chow, T.Y., Lin, T.P., Hsing, Y.I.: Gene cloning and characterization of a soybean (*Glycine max* L.) LEA protein, GmPM16. - *Plant mol. Biol.* **56**: 689-703, 2004.
- Shinozaki, K., Yamaguchi-Shinozaki, K.: Molecular response to dehydration and low temperature: differences and cross-talk between two stress signaling pathways. - *Curr. Opin. Plant Biol.* **3**: 217-223, 2000.
- Shou, H., Bordallo, P., Wang, K.: Expression of the *Nicotiana* protein kinase (NPK1) enhanced drought tolerance in transgenic maize. - *J. exp. Bot.* **55**: 1013-1019, 2004.
- Shridhar, R.J., Vadez, V., Mathur, P.B., Narasu, M.L., Sharma, K.K.: Better root:shoot ratio conferred enhanced harvest index in transgenic groundnut overexpressing the *rd29A:DREB1A* gene under intermittent drought stress in an outdoor lysimetric dry-down trial. - *J. semi-arid tropics Agr. Res.* **10**: 1-7, 2012.
- Siefert, F., Tyree, M. T., Lovisolo, C., Schubert, A., Kaldenhoff, R.: PIP1 plasma membrane aquaporins in tobacco: from cellular effects to function in plants. - *Plant Cell* **14**: 869-876, 2002.
- Singh, N., Mishra, A., Jha, B.: Over-expression of the peroxisomal ascorbate peroxidase (*SbpAPX*) gene cloned from halophyte *Salicornia brachiata* confers salt and drought stress tolerance in transgenic tobacco. - *Mar. Biotechnol.* **16**: 321-332, 2014.
- Singh, N.K., Nelson, D.E., Kuhn, D., Hasegawa, P.M., Bressan, R.A.: Molecular cloning of osmotin and regulation of its expression by ABA and adaptation to low water potential. - *Plant Physiol.* **90**: 1096-1101, 1989.
- Sivamani, E., Bahieldin, A., Wraith, J.M., Al-Niemi, T., Dyer, W.E., Ho, T.H.D., Qu, R.D.: Improved biomass productivity and water use efficiency under water deficit conditions in transgenic wheat constitutively expressing the barley HVA1 gene. - *Plant Sci.* **155**: 1-9, 2000.
- Skirycz, A., Memmi, S., De Bodt, S., Maleux, K., Obata, T., Fernie, A.R., Devreese, B., Inze, D.: A reciprocal 15N-labeling proteomic analysis of expanding *Arabidopsis* leaves subjected to osmotic stress indicates importance of mitochondria in preserving plastid functions. - *J. Proteome Res.* **10**: 1018-1029, 2011.
- Smirnoff, N., Wheeler, G.L.: Ascorbic acid in plants: biosynthesis and function. - *CRC Crit. Rev. Plant. Sci.* **19**: 267-290, 2000.
- Spollen, W.G., Nelson, C.J.: Response of fructan to water-deficit in growing leaves of tall fescue. - *Plant Physiol.* **106**: 329-336, 1994.
- Stiller, I., Dulai, S., Kondrák, M., Tarnai, R., Szabó, L., Toldi, O., Bánfalvi, Z.: Effects of drought on water content and photosynthetic parameters in potato plants expressing the trehalose-6-phosphate synthase gene of *Saccharomyces cerevisiae*. - *Planta* **227**: 299-308, 2008.
- Sujata, B., Kshitija, S.: Drought stress adaptation: metabolic adjustment and regulation of gene expression. - *Plant Breed.* **132**: 21-32, 2013.
- Sun, W., Bernard, C., Van de Cotte, B., Van Montagu, M., Verbruggen, N.: *At-HSP17.6A*, encoding a small heat-shock protein in *Arabidopsis*, can enhance osmotolerance upon overexpression. - *Plant J.* **27**: 407-415, 2001.
- Sun, W.H., Duan, M., Shu, D.F., Yang, S., Meng, Q.W.: Over-expression of *StAPX* in tobacco improves seed germination and increases early seedling tolerance to salinity and osmotic stresses. - *Plant Cell Rep.* **29**: 917-926, 2010.
- Šurbanovski, N., Sargent, D.J., Else, M.A., Simpson, D.W., Zhang, H., Grant, O.M.: Expression of *Fragaria vesca* PIP aquaporins in response to drought stress: PIP down-regulation correlates with the decline in substrate moisture content. - *PLoS ONE* **8**: e74945, 2013.
- Szabados, L., Savoure, A.: Proline: a multifunctional amino acid. - *Trends Plant Sci.* **15**: 89-97, 2010.
- Székely, G., Abrahám, E., Cséplő, A., Rigó, G., Zsigmond, L., Csiszár, J., Ayaydin, F., Strizhov, N., Jásik, J., Schmelzer, E., Koncz, C., Szabados, L.: Duplicated *P5CS* genes of *Arabidopsis* play distinct roles in stress regulation and developmental control of proline biosynthesis. - *Plant J.* **53**: 11-28, 2008.
- Tada, Y., Spoel, S.H., Pajeroska-Mukhtar, K., Mou, Z., Song, J., Wang, C., Zuo, J., Dong, X.: Plant immunity requires conformational changes of NPR1 via nitrosylation and thioredoxins. - *Science* **321**: 952-956, 2008.
- Taji, T., Seki, M., Satou, M., Sakurai, T., Kobayashi, M., Ishiyama, K., Narusaka, Y., Narusaka, M., Zhu, J.K., Shinozaki, K.: Comparative genomics in salt tolerance between *Arabidopsis* and *Arabidopsis*-related halophyte salt cress using *Arabidopsis* microarray. - *Plant. Physiol.* **135**: 1697-1709, 2004.
- Tarczynski, M.C., Jensen, R.G., Bohnert, H.J.: Expression of a bacterial *mtlD* gene in transgenic tobacco leads to production and accumulation of mannitol. - *Proc. nat. Acad. Sci. USA.* **89**: 2600-2604, 1992.
- Tarczynski, M.C., Jensen, R.G., Bohnert, H.J.: Stress protection of transgenic tobacco by production of the osmolyte mannitol. - *Science* **259**: 508-510, 1993.
- Taylor, L., Nunes-Nesi, A., Parsley, K., Leiss, A., Leach, G., Coates, S., Wingler, A., Fernie, A.R., Hibberd, J.M.: Cytosolic pyruvate, orthophosphate dikinase functions in nitrogen remobilization during leaf senescence and limits individual seed growth and nitrogen content. - *Plant J.* **62**: 641-652, 2010.
- Thomas, J.C., Sepahi, M., Arendall, B., Bohnert, H.J.: Enhancement of seed germination in high salinity by engineering mannitol expression in *Arabidopsis thaliana*. - *Plant Cell Environ.* **18**: 801-806, 1995.
- Trujillo, L.E., Menéndez, C., Ochogavía, M.E., Hernández, I., Borrás, O., Rodríguez, R., Coll, Y., Arrieta, J.G., Banguela, A., Ramírez, R., Hernández, L.: Engineering drought and salt tolerance in plants using *SodERF3*, a novel sugarcane ethylene responsive factor. - *Biotechnol. Appl.* **2**: 168-171, 2009.
- Uga, Y., Sugimoto, K., Ogawa, S., Rane, J., Ishitani, M., Hara, N., Kitomi, Y., Inukai, Y., Ono, K., Kanno, N., Inoue, H., Takehisa, H., Motoyama, R., Nagamura, Y., Wu, J., Matsumoto, T., Takai, T., Okuno, K., Yano, M.: Control of root system architecture by *DEEPER ROOTING 1* increases rice yield under drought conditions. - *Nat. Genet.* **45**: 1097-1102, 2013.
- Ulker, B., Somssich, I.E.: WRKY transcription factors: from DNA binding towards biological function. - *Curr. Opin. Plant Biol.* **7**: 491-498, 2004.
- Uno, Y., Furihata, T., Abe, H., Yoshida, R., Shinozaki, K., Yamaguchi-Shinozaki, K.: *Arabidopsis* basic leucine zipper transcriptional factors involved in an abscisic acid-dependent signal transduction pathway under drought and high-salinity conditions. - *Proc. nat. Acad. Sci. USA* **97**: 11632-11637, 2000.
- Vaahtera, L., Brosché, M.: More than the sum of its parts - how to achieve a specific transcriptional response to abiotic stress. - *Plant Sci.* **180**: 421-430, 2011.

- Valluru, R., Van den Ende, W.: Plant fructans in stress environments: emerging concepts and future prospects. - J. exp. Bot. **59**: 2905-2916, 2008.
- Van den Boogaard, Alewijnse, D., Veneklaas E.J., Lambers, H.: Growth and water-use efficiency of ten *Triticum aestivum* cultivars at different water availability in relation to allocation of biomass. - Plant Cell Environ. **20**: 200-210, 1997.
- Vellosillo, T., Vicente, J., Kulasekaran, S., Hamberg, M., Castresana, C.: Emerging complexity in reactive oxygen species production and signaling during the response of plants to pathogens. - Plant Physiol. **154**: 444-448, 2010.
- Venkataraman, S., Schafer, F.Q., Buettner, G.: Detection of lipid radicals using EPR. - Antioxidants Redox Signal. **6**: 631-638, 2004.
- Verelst, W., Bertolini, E., De Bodt, S., Vandepoele, K., Demeulenaere, M., Pe, M.E., Inze, D.: Molecular and physiological analysis of growth-limiting drought stress in *Brachypodium distachyon* leaves. - Mol. Plant. **6**: 311-322, 2013.
- Versulues, P.E., Agarwal, M., Katiyar-Agarwal, S., Zhu, J., Zhu, J.K.: Methods and concepts in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status. - Plant J. **45**: 523-539, 2006.
- Vinocur, B., Altman, A.: Recent advances in engineering plant tolerance to abiotic stress: achievements and limitations. - Curr. Opin. Biotechnol. **16**: 123-32, 2005.
- Wang, P., Du, Y., Zhao, X., Miao, Y., Song, C.P.: The MPK6-ERF6-ROSE7/GCC-box complex modulates oxidative gene transcription and the oxidative response in *Arabidopsis thaliana*. - Plant Physiol. **161**: 1392-1408, 2013.
- Wang, P., Duan, W., Takabayashi, A., Endo, T., Shikanai, T., Ji-Yu Ye., Hualing, M.: Chloroplastic NAD(P)H dehydrogenase in tobacco leaves functions in alleviation of oxidative damage caused by temperature stress. - Plant Physiol. **141**: 465-474, 2006.
- Witcombe, J.R., Hollington, P.A., Howarth, C.J., Reader, S., Steele, K.A.: Breeding for abiotic stresses for sustainable agriculture. - Phil. Trans. roy. Soc. London B **363**: 703-716, 2008.
- Wrzaczek, M., Vainonen, J.P., Gauthier, A., Overmyer, K., Kangasjärvi, J.: Reactive oxygen in abiotic stress perception - from genes to proteins. - In: Shanker, A., Venkateswarlu, B. (ed.): Abiotic Stress Response in Plants - Physiological, Biochemical and Genetic Perspectives. Pp. 27-40. In-Tech, 2011.
- Wu, G., Wei, Z.K., Shao, H.B.: The mutual responses of higher plants to environment: physiological and microbiological aspects. - Biointerface **59**: 113-119, 2007.
- Wu, X., Shiroto, Y., Kishitani, S., Ito, Y., Toriyama, K.: Enhanced heat and drought tolerance in transgenic rice seedlings overexpressing *OsWRKY11* under the control of HSP101 promoter. - Plant Cell Rep. **28**: 21-30, 2009.
- Xiao, B., Huang, Y., Tang, N., Xiong, L.: Over-expression of a *LEA* gene in rice improves drought resistance under the field conditions. - Theor. appl. Genet. **115**: 35-46, 2007.
- Xoconostle-Cazares, B., Ramirez-Ortega, F.A., Flores-Elenes, L., Ruiz-Medrano, R.: Drought tolerance in crop plants. - Amer. J. Plant Physiol. **5**: 241-256, 2011.
- Xu, D., Duan, X., Wang, B., Hong, B., Ho, T., Wu, R.: Expression of a late embryogenesis abundant protein gene, *HVA1*, from barley confers tolerance to water deficit and salt stress in transgenic rice. - Plant Physiol. **110**: 249-257, 1996.
- Xu, D.Q., Huang, J., Guo, S.Q., Yang, X., Bao, Y.-M., Tang, H.-J., Zhang, H.-S.: Overexpression of a TFIIIA-type zinc finger protein gene *ZFP252* enhances drought and salt tolerance in rice (*Oryza sativa* L.). - FEBS Lett. **582**: 1037-1043, 2008.
- Xu, Y., Hu, W., Liu, J., Zhang, J., Jia, C., Miao, H., Jin, B.X.Z.: A banana aquaporin gene, *MaPIPI1*, is involved in tolerance to drought and salt stresses. - BMC Plant Biol. **14**: 59, 2014.
- Yamaguchi-Shinozaki, K., Shinozaki, K.: Organization of *cis*-acting regulatory elements in osmotic- and cold-stress-responsive promoters. - Trends Plant Sci. **10**: 88-94, 2005.
- Yancey, P.H., Clark, M.E., Hand, S.C., Bowlus, R.D., Somero, G.N.: Living with water stress: evolution of osmolyte systems. - Science **217**: 1214-1222, 1982.
- Yang, J., Zhang, J., Wang, Z., Zhu, Q., Liu, L.: Absciscic acid and cytokinins in the root exudates and leaves and their relationship to senescence and remobilization of carbon reserves in rice subjected to water stress during grain filling. - Planta **215**: 645-652, 2002.
- Yang, W.J., Rich, P.J., Axtell, J.D., Wood, K.V., Bonham, C.C., Ejeta, G., Mickelbart, M.V., Rhodes, D.: Genotypic variation for glycine betaine in sorghum. - Crop Sci. **43**: 162-169, 2003.
- Yordanov, I., Velikova, V., Tsonev, T.: Plant responses to drought and stress tolerance. - Bulg. J. Plant Physiol. **2003** (Special issue): 187-206, 2003.
- Yoshida, T., Mogami, J., Yamaguchi-Shinozaki, K.: Omics approaches toward defining the comprehensive abscisic acid signaling network in plants. - Plant Cell Physiol. **56**: 1043-1052, 2015.
- Zhang, L., Zhao, G., Xia, C., Jia, J., Liu, X., Kong, X.: A wheat R2R3-MYB gene, *TaMYB30-B*, improves drought stress tolerance in transgenic *Arabidopsis*. - J. exp. Bot. **63**: 5873-5885, 2012.
- Zhang, L., Tian, L.H., Zhao, J.F., Song, Y., Zhang, C.J., Guo, Y.: Identification of an apoplastic protein involved in the initial phase of salt stress response in rice root by two-dimensional electrophoresis. - Plant Physiol. **149**: 916-928, 2009.
- Zhang, X., Wollenweber, B., Jiang, D., Liu, F., Zhao, J.: Water deficits and heat shock effects on photosynthesis of a transgenic *Arabidopsis thaliana* constitutively expressing ABP9, a bZIP transcription factor. - J. exp. Bot. **59**: 839-848, 2008a.
- Zhang, Y., Yang, J., Lu, S., Cai, J., Guo, Z.: Overexpressing *SgNCED1* in tobacco increases ABA level, antioxidant enzyme activities, and stress tolerance. - J. Plant Growth Regul. **27**: 151-158, 2008b.
- Zhou, S., Hu, W., Deng, X., Ma, Z., Chen, L., Huang, C., Wang, C., Wang, J., He, Y., Yang, G., He, G.: Over-expression of the wheat aquaporin gene, *TaAQP7*, enhances drought tolerance in transgenic tobacco. - PLoS ONE **7**: e52439, 2012.
- Zhu, J.K.: Salt and drought stress signal transduction in plants. - Annu. Rev. Plant Biol. **53**: 247-273, 2002.
- Zlatev, Z., Lidon, F.C.: An overview on drought induced changes in plant growth, water relations and photosynthesis. - Emir. J. Food Agr. **24**: 57-72, 2012.